

AL  
PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : A61K 45/06, 31/585, 31/41	A2	(11) International Publication Number: WO 96/40258
		(43) International Publication Date: 19 December 1996 (19.12.96)

(21) International Application Number: PCT/US96/09342  
(22) International Filing Date: 5 June 1996 (05.06.96)  
(30) Priority Data:  
08/486,089 7 June 1995 (07.06.95) US  
(60) Parent Application or Grant  
(63) Related by Continuation  
US 08/486,089 (CON)  
Filed on 7 June 1996 (07.06.96)  
(71) Applicant (for all designated States except US): G.D. SEARLE  
& CO. [US/US]; Corporate Patent Dept., P.O. Box 5110,  
Chicago, IL 60680-5110 (US).  
(72) Inventors; and  
(75) Inventors/Applicants (for US only): MACLAUGHLAN, Todd,  
E. [CA/US]; 34128 Sulkey Drive, Grayslake, IL 60093  
(US). SCHUH, Joseph, R. [US/US]; 2055 Rurline Drive,  
St. Louis, MO 63146 (US).  
(74) Agents: KEANE, J., Timothy et al.; G.D. Searle & Co.,  
Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-  
5110 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY,  
CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL,  
IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV,  
MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU,  
SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ,  
VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT,  
LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI,  
CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

*Without international search report and to be republished  
upon receipt of that report.*

(54) Title: SPIRONOLACTONE AND ANGIOTENSIN II ANTAGONIST COMBINATION THERAPY FOR TREATMENT OF  
CONGESTIVE HEART FAILURE

(57) Abstract

A combination therapy comprising a therapeutically-effective amount of an epoxy-free spiro lactone-type aldosterone receptor antagonist and a therapeutically-effective amount of an angiotensin II receptor antagonist is described for treatment of circulatory disorders, including cardiovascular disorders such as hypertension, congestive heart failure, cirrhosis and ascites. Preferred angiotensin II receptor antagonists are those compounds having high potency and bioavailability and which are characterized in having an imidazole or triazole moiety attached to a biphenylmethyl or pyridinyl/phenylmethyl moiety. A preferred epoxy-free spiro lactone-type aldosterone receptor antagonist is spironolactone. A preferred combination therapy includes the angiotensin II receptor antagonist 5-[2-{5-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl]-2-pyridinyl}phenyl-1H-tetrazole and the aldosterone receptor antagonist spironolactone.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

SPIRONOLACTONE AND ANGIOTENSIN II ANTAGONIST  
COMBINATION THERAPY FOR TREATMENT OF CONGESTIVE  
HEART FAILURE

5

Field of the Invention

Combinations of a spiro lactone-type aldosterone  
receptor antagonist and an angiotensin II receptor  
10 antagonist are described for use in treatment of  
circulatory disorders, including cardiovascular diseases  
such as hypertension, congestive heart failure, cirrhosis  
and ascites. Of particular interest are therapies using  
an epoxy-free spiro lactone-type aldosterone receptor  
15 antagonist compound such as spironolactone in combination  
with an angiotensin II receptor antagonist compound.

Background of the Invention

20 Myocardial (or cardiac) failure, whether a  
consequence of a previous myocardial infarction, heart  
disease associated with hypertension, or primary  
cardiomyopathy, is a major health problem of worldwide  
proportions. The incidence of symptomatic heart failure  
25 has risen steadily over the past several decades.

In clinical terms, decompensated cardiac  
failure consists of a constellation of signs and symptoms  
that arises from congested organs and hypoperfused  
30 tissues to form the congestive heart failure (CHF)  
syndrome. Congestion is caused largely by increased  
venous pressure and by inadequate sodium ( $\text{Na}^+$ ) excretion,  
relative to dietary  $\text{Na}^+$  intake, and is importantly  
related to circulating levels of aldosterone (ALDO). An  
35 abnormal retention of  $\text{Na}^+$  occurs via tubular epithelial  
cells throughout the nephron, including the later portion  
of the distal tubule and cortical collecting ducts, where

ALDO receptor sites are present.

ALDO is the body's most potent mineralocorticoid hormone. As connoted by the term  
5 mineralocorticoid, this steroid hormone has mineral-regulating activity. It promotes  $\text{Na}^+$  reabsorption not only in the kidney, but also from the lower gastrointestinal tract and salivary and sweat glands, each of which represents classic ALDO-responsive tissues.  
10 ALDO regulates  $\text{Na}^+$  and water resorption at the expense of potassium ( $\text{K}^+$ ) and magnesium ( $\text{Mg}^{2+}$ ) excretion.

ALDO can also provoke responses in nonepithelial cells. Elicited by a chronic elevation in  
15 plasma ALDO level that is inappropriate relative to dietary  $\text{Na}^+$  intake, these responses can have adverse consequences on the structure of the cardiovascular system. Hence, ALDO can contribute to the progressive nature of myocardial failure for multiple reasons.

20 Multiple factors regulate ALDO synthesis and metabolism, many of which are operative in the patient with myocardial failure. These include renin as well as non-renin-dependent factors (such as  $\text{K}^+$ , ACTH) that  
25 promote ALDO synthesis. Hepatic blood flow, by regulating the clearance of circulating ALDO, helps determine its plasma concentration, an important factor in heart failure characterized by reduction in cardiac output and hepatic blood flow.

30 The renin-angiotensin-aldosterone system (RAAS) is one of the hormonal mechanisms involved in regulating pressure/volume homeostasis and also in the development of hypertension. Activation of the renin-angiotensin-aldosterone system begins with renin secretion from the  
35 juxtaglomerular cells in the kidney and culminates in the formation of angiotensin II, the primary active species

of this system. This octapeptide, angiotensin II, is a potent vasoconstrictor and also produces other physiological effects such as stimulating aldosterone secretion, promoting sodium and fluid retention, inhibiting renin secretion, increasing sympathetic nervous system activity, stimulating vasopressin secretion, causing positive cardiac inotropic effect and modulating other hormonal systems.

Previous studies have shown that antagonizing angiotensin II binding at its receptors is a viable approach to inhibit the renin-angiotensin system, given the pivotal role of this octapeptide which mediates the actions of the renin-angiotensin system through interaction with various tissue receptors. There are several known angiotensin II antagonists, most of which are peptidic in nature. Such peptidic compounds are of limited use due to their lack of oral bioavailability or their short duration of action. Also, commercially-available peptidic angiotensin II antagonists (e.g., Saralasin) have a significant residual agonist activity which further limit their therapeutic application.

Non-peptidic compounds with angiotensin II antagonist properties are known. For example, early descriptions of such non-peptidic compounds include the sodium salt of 2-n-butyl-4-chloro-1-(2-chlorobenzyl)imidazole-5-acetic acid which has specific competitive angiotensin II antagonist activity as shown in a series of binding experiments, functional assays and in vivo tests [P. C. Wong et al, J. Pharmacol. Exp. Ther., 247(1), 1-7 (1988)]. Also, the sodium salt of 2-butyl-4-chloro-1-(2-nitrobenzyl)imidazole-5-acetic acid has specific competitive angiotensin II antagonist activity as shown in a series of binding experiments, functional assays and in vivo tests [A. T. Chiu et al, European J. Pharmacol., 157, 31-21 (1988)]. A family of

1-benzylimidazole-5-acetate derivatives has been shown to have competitive angiotensin II antagonist properties [A. T. Chiu et al, J. Pharmacol. Exp. Ther., 250(3), 867-874 (1989)]. U.S. Patent No. 4,816,463 to Blankey et al describes a family of 4,5,6,7-tetrahydro-1H-imidazo(4,5-c)-tetrahydro-pyridine derivatives useful as antihypertensives, some of which are reported to antagonize the binding of labelled angiotensin II to rat adrenal receptor preparation and thus cause a significant decrease in mean arterial blood pressure in conscious hypertensive rats. Other families of non-peptidic angiotensin II antagonists have been characterized by molecules having a biphenylmethyl moiety attached to a heterocyclic moiety. For example, EP No. 253,310, published 20 January 1988, describes a series of aralkyl imidazole compounds, including in particular a family of biphenylmethyl substituted imidazoles, as antagonists to the angiotensin II receptor. EP No. 323,841 published 12 July 1989 describes four classes of angiotensin II antagonists, namely, biphenylmethylpyrroles, biphenylmethylpyrazoles, biphenylmethyl-1,2,3-triazoles and biphenylmethyl 4-substituted-4H-1,2,4-triazoles, including the compound 3,5-dibutyl-4-[(2'-carboxybiphenyl-4-yl)methyl]-4H-1,2,4-triazole. U.S. Patent No. 4,880,804 to Carini et al describes a family of biphenylmethylbenzimidazole compounds as angiotensin II receptor blockers for use in treatment of hypertension and congestive heart failure.

Many aldosterone receptor blocking drugs are known. For example, spironolactone is a drug which acts at the mineralocorticoid receptor level by competitively inhibiting aldosterone binding. This steroidal compound has been used for blocking aldosterone-dependent sodium transport in the distal tubule of the kidney in order to reduce edema and to treat essential hypertension and primary hyperaldosteronism [F. Mantero et al, Clin. Sci.

Mol. Med., 45 (Suppl 1), 219s-224s (1973)].

Spironolactone is also used commonly in the treatment of other hyperaldosterone-related diseases such as liver cirrhosis and congestive heart failure [F.J. Saunders et al, Aldactone; Spironolactone: A Comprehensive Review, Searle, New York (1978)]. Progressively-increasing doses of spironolactone from 1 mg to 400 mg per day [i.e., 1 mg/day, 5 mg/day, 20 mg/day] were administered to a spironolactone-intolerant patient to treat cirrhosis-related ascites [P.A. Greenberger et al, N. Eng. J. Med., 314(4), 343-345 (Jul-Aug, 1986)]. It has been recognized that development of myocardial fibrosis is sensitive to circulating levels of both Angiotensin II and aldosterone, and that the aldosterone antagonist spironolactone prevents myocardial fibrosis in animal models, thereby linking aldosterone to excessive collagen deposition [D. Klug et al, Am. J. Cardiol., 71 (3), 46A-54A (1993)]. Spironolactone has been shown to prevent fibrosis in animal models irrespective of the development of left ventricular hypertrophy and the presence of hypertension [C.G. Brilla et al, J. Mol. Cell. Cardiol., 25(5), 563-575 (1993)]. Spironolactone at a dosage ranging from 25 mg to 100 mg daily is used to treat diuretic-induced hypokalemia, when orally-administered potassium supplements or other potassium-sparing regimens are considered inappropriate [Physicians' Desk Reference, 46th Edn., p. 2153, Medical Economics Company Inc., Montvale, N.J. (1992)].

Previous studies have shown that inhibiting ACE inhibits the renin-angiotensin system by substantially complete blockade of the formation of angiotensin II. Many ACE inhibitors have been used clinically to control hypertension. While ACE inhibitors may effectively control hypertension, side effects are common including chronic cough, skin rash, loss of taste sense, proteinuria and neutropenia.

Moreover, although ACE inhibitors effectively block the formation of angiotensin II, aldosterone levels are not well controlled in certain patients having cardiovascular diseases. For example, despite continued ACE inhibition in hypertensive patients receiving captopril, there has been observed a gradual return of plasma aldosterone to baseline levels [J. Staessen et al, J. Endocrinol., 91, 457-465 (1981)]. A similar effect has been observed for patients with myocardial infarction receiving zofenopril [C. Borghi et al, J. Clin. Pharmacol., 33, 40-45 (1993)]. This phenomenon has been termed "aldosterone escape".

Another series of steroidal-type aldosterone receptor antagonists is exemplified by epoxy-containing spironolactone derivatives. For example, U.S. Patent No. 4,559,332 issued to Grob et al describes 9 $\alpha$ ,11 $\alpha$ -epoxy-containing spironolactone derivatives as aldosterone antagonists useful as diuretics. These 9 $\alpha$ ,11 $\alpha$ -epoxy steroids have been evaluated for endocrine effects in comparison to spironolactone [M. de Gasparo et al, J. Pharm. Exp. Ther., 240(2), 650-656 (1987)].

Combinations of an aldosterone antagonist and an ACE inhibitor have been investigated for treatment of heart failure. It is known that mortality is higher in patients with elevated levels of plasma aldosterone and that aldosterone levels increase as CHF progresses from activation of the Renin-Angiotensin-Aldosterone System (RAAS). Routine use of a diuretic may further elevate aldosterone levels. ACE inhibitors consistently inhibit angiotensin II production but exert only a mild and transient antialdosterone effect.

35

Combining an ACE inhibitor and spironolactone has been suggested to provide substantial inhibition of the



entire RAAS. For example, a combination of enalapril and spironolactone has been administered to ambulatory patients with monitoring of blood pressure [P. Poncelet et al, Am. J. Cardiol., 65(2), 33K-35K (1990)]. In a 90-  
5 patient study, a combination of captopril and spironolactone was administered and found effective to control refractory CHF without serious incidents of hyperkalemia [U. Dahlstrom et al, Am. J. Cardiol., 71, 29A-33A (21 Jan 1993)]. Spironolactone coadministered  
10 with an ACE inhibitor was reported to be highly effective in 13 of 16 patients afflicted with congestive heart failure [A.A. van Vliet et al, Am. J. Cardiol., 71, 21A-28A (21 Jan 1993)]. Clinical improvements have been reported for patients receiving a co-therapy of  
15 spironolactone and the ACE inhibitor enalapril, although this report mentions that controlled trials are needed to determine the lowest effective doses and to identify which patients would benefit most from combined therapy [F. Zannad, Am. J. Cardiol., 71(3), 34A-39A (1993)].

20

Combinations of an angiotensin II receptor antagonist and aldosterone receptor antagonist, are known. For example, PCT Application No. US91/09362 published 25 June 1992 describes treatment of  
25 hypertension using a combination of an imidazole-containing angiotensin II antagonist compound and a diuretic such as spironolactone.

Summary of the Invention

A combination therapy comprising a therapeutically-effective amount of an angiotensin II  
5 receptor antagonist and a therapeutically-effective amount of an epoxy-free spirolactone-type aldosterone receptor antagonist is useful to treat circulatory disorders, including cardiovascular disorders such as hypertension, congestive heart failure, cirrhosis and  
10 ascites.

The phrase "angiotensin II receptor antagonist" is intended to embrace one or more compounds or agents having the ability to interact with a receptor site  
15 located on various human body tissues, which site is a receptor having a relatively high affinity for angiotensin II and which receptor site is associated with mediating one or more biological functions or events such as vasoconstriction or vasorelaxation, kidney-mediated  
20 sodium and fluid retention, sympathetic nervous system activity, and in modulating secretion of various substances such as aldosterone, vasopressin and renin, to lower blood pressure in a subject susceptible to or afflicted with elevated blood pressure. Interactions of  
25 such angiotensin II receptor antagonist with this receptor site may be characterized as being either "competitive" (i.e., "surmountable") or as being "insurmountable". These terms, "competitive" and "insurmountable", characterize the relative rates, faster  
30 for the former term and slower for the latter term, at which the antagonist compound dissociates from binding with the receptor site.

The phrase "epoxy-free spirolactone-type  
35 aldosterone receptor antagonist" embraces an agent or compound, or a combination of two or more of such agents or compounds, which agent or compound binds to the

aldosterone receptor as a competitive inhibitor of the action of aldosterone itself at the receptor site in the renal tubules, so as to modulate the receptor-mediated activity of aldosterone. Typical of such aldosterone  
5 receptor antagonists are spiro lactone-type compounds. The term "spiro lactone-type" is intended to characterize a steroidal structure comprising a lactone moiety attached to a steroid nucleus, typically at the steroid "D" ring, through a spiro bond configuration. Preferred  
10 spiro lactone-type compounds are epoxy-free, e.g., compounds which do not contain an epoxy moiety attached to any portion of the steroid nucleus.

The phrase "combination therapy", in defining  
15 use of an angiotensin II antagonist and a spiro lactone-type aldosterone receptor antagonist, is intended to embrace administration of each antagonist in a sequential manner in a regimen that will provide beneficial effects of the drug combination, and is intended to embrace co-  
20 administration of the antagonist agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of active ingredients or in multiple, separate capsules for each antagonist agent.

25 The phrase "therapeutically-effective" is intended to qualify the amount of each antagonist agent for use in the combination therapy which will achieve the goal of reduction of hypertension with improvement in cardiac sufficiency by reducing or preventing, for  
30 example, hypertension and/or the progression of congestive heart failure.

The phrase "low-dose amount", in characterizing a therapeutically-effective amount of the aldosterone  
35 receptor antagonist agent in the combination therapy, is intended to define a quantity of such agent, or a range of quantity of such agent, that is capable of improving

cardiac sufficiency while reducing or avoiding one or more aldosterone-antagonist-induced side effects, such as hyperkalemia. A dosage of an aldosterone receptor antagonist, e.g., spironolactone, which would accomplish  
5 the therapeutic goal of favorably enhancing cardiac sufficiency, while reducing or avoiding side effects, would be a dosage that substantially avoids inducing diuresis, that is, a substantially non-diuresis-effective dosage or a non-diuretic-effective amount of an  
10 aldosterone receptor antagonist.

Another combination therapy of interest would consist essentially of three active agents, namely, an AII antagonist, an aldosterone receptor antagonist agent  
15 and a diuretic.

For a combination of AII antagonist agent and an ALDO antagonist agent, the agents would be used in combination in a weight ratio range from about 0.5-to-one  
20 to about twenty-to-one of the AII antagonist agent to the aldosterone receptor antagonist agent. A preferred range of these two agents (AII antagonist-to-ALDO antagonist) would be from about one-to-one to about fifteen-to-one, while a more preferred range would be from about one-to-  
25 one to about five-to-one, depending ultimately on the selection of the AII antagonist and ALDO antagonist. The diuretic agent may be present in a ratio range of 0.1-to-one to about ten to one (AII antagonist to diuretic).

Detailed Description of the Invention

Examples of angiotensin II (AII) antagonists which may be used in the combination therapy are shown in the following categories:

A first group of AII antagonists consists of the following compounds:

- saralasin acetate, candesartan cilexetil, CGP-63170,  
10 EMD-66397, KT3-671, LR-B/081, valsartan, A-81282,  
BIBR-363, BIBS-222, BMS-184698, candesartan, CV-11194,  
EXP-3174, KW-3433, L-161177, L-162154, LR-B/057,  
LY-235656, PD-150304, U-96849, U-97018, UP-275-22,  
WAY-126227, WK-1492.2K, YM-31472, losartan potassium,  
15 E-4177, EMD-73495, eprosartan, HN-65021, irbesartan,  
L-159282, ME-3221, SL-91.0102, Tasosartan, Telmisartan,  
UP-269-6, YM-358, CGP-49870, GA-0056, L-159689, L-162234,  
L-162441, L-163007, PD-123177, A-81988, BMS-180560,  
CGP-38560A, CGP-48369, DA-2079, DE-3489, DuP-167,  
20 EXP-063, EXP-6155, EXP-6803, EXP-7711, EXP-9270, FK-739,  
HR-720, ICI-D6888, ICI-D7155, ICI-D8731, isoteoline,  
KRI-1177, L-158809, L-158978, L-159874, LR B087,  
LY-285434, LY-302289, LY-315995, RG-13647, RWJ-38970,  
RWJ-46458, S-8307, S-8308, saprisartan, saralasin,  
25 Sarmesin, WK-1360, X-6803, ZD-6888, ZD-7155, ZD-8731,  
BIBS39, CI-996, DMP-811, DuP-532, EXP-929, L-163017,  
LY-301875, XH-148, XR-510, zolasartan and PD-123319.

A second group of AII antagonists of interest consists of the following compounds:

- saralasin acetate, candesartan cilexetil, CGP-63170,  
EMD-66397, KT3-671, LR-B/081, valsartan, A-81282,  
BIBR-363, BIBS-222, BMS-184698, candesartan, CV-11194,  
EXP-3174, KW-3433, L-161177, L-162154, LR-B/057,  
35 LY-235656, PD-150304, U-96849, U-97018, UP-275-22,  
WAY-126227, WK-1492.2K, YM-31472, losartan potassium,  
E-4177, EMD-73495, eprosartan, HN-65021, irbesartan,

12

L-159282, ME-3221, SL-91.0102, Tasosartan, Telmisartan, UP-269-6, YM-358, CGP-49870, GA-0056, L-159689, L-162234, L-162441, L-163007 and PD-123177.

5           A family of spirolactone-type compounds of interest for use in the combination therapy is defined by Formula A

10

(A)

          wherein R is lower alkyl of up to 5 carbon  
15 atoms, and

20           Lower alkyl residues include branched and unbranched groups, preferably methyl, ethyl and n-propyl.

Specific compounds of interest within Formula A are the following:

25           7 $\alpha$ -Aceylthio-3-oxo-4,15-androstadiene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
          3-Oxo-7 $\alpha$ -propionylthio-4,15-androstadiene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
          6 $\beta$ ,7 $\beta$ -Methylene-3-oxo-4,15-androstadiene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
30           15 $\alpha$ ,16 $\alpha$ -Methylene-3-oxo-4,7 $\alpha$ -propionylthio-4-androstene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
          6 $\beta$ ,7 $\beta$ ,15 $\alpha$ ,16 $\alpha$ -Dimethylene-3-oxo-4-androstene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
35           7 $\alpha$ -Aceylthio-15 $\beta$ ,16 $\beta$ -Methylene-3-oxo-4-androstene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
          15 $\beta$ ,16 $\beta$ -Methylene-3-oxo-7 $\beta$ -propionylthio-4-

androstene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one; and  
6 $\beta$ ,7 $\beta$ ,15 $\beta$ ,16 $\beta$ -Dimethylene-3-oxo-4-androstene-[17( $\beta$ -  
1')-spiro-5']perhydrofuran-2'-one.

- 5 Methods to make compounds of Formula A are described in  
U.S. Patent No. 4,129,564 to Wiechart et al issued on 12  
December 1978.

A second family of spirolactone-type compounds of  
10 interest for use in the combination therapy is defined by  
Formula B:

15

(B)

wherein

- 20  $R^1$  is  $C_{1-3}$ -alkyl or  $C_{1-3}$  acyl and  $R^2$  is hydrogen or  $C_{1-3}$ -  
alkyl.

Specific compounds of interest within Formula B are the  
following:

- 25 1 $\alpha$ -Acetylthio-15 $\beta$ ,16 $\beta$ -methylene-7 $\alpha$ -methylthio-3-oxo-  
17 $\alpha$ -pregn-4-ene-21,17-carbolactone; and  
15 $\beta$ ,16 $\beta$ -Methylene-1 $\alpha$ ,7 $\alpha$ -dimethylthio-3-oxo-17 $\alpha$ -  
pregn-4-ene-21,17-carbolactone.

- 30 Methods to make the compounds of Formula B are described  
in U.S. Patent No. 4,789,668 to Nickisch et al which  
issued 6 December 1988.

A third family of spirolactone-type compounds  
35 of interest for use in the combination therapy is defined  
by a structure of Formula C:

5

(C)

Specific compounds of interest include:

7 $\alpha$ -Acylthio-21-hydroxy-3-oxo-17 $\alpha$ -pregn-4-ene-17-carboxylic acid lactones; and

10

21-hydroxy-3-oxo-17 $\alpha$ -pregn-1,4-diene-17-carboxylic acid lactone.

Methods to make the compounds of Formula C are described in U.S. Patent No. 3,257,390 to Patchett which issued 21 June 1966. Of particular interest is the compound spironolactone having the following structure and formal name:

20

25 "spironolactone": 17-hydroxy-7 $\alpha$ -mercapto-3-oxo-17 $\alpha$ -pregn-4-ene-21-carboxylic acid  $\gamma$ -lactone acetate


Spironolactone is sold by G.D. Searle & Co., Skokie, Illinois, under the trademark "ALDACTONE", in tablet dosage form at doses of 25 mg, 50 mg and 100 mg per tablet.

A diuretic agent may be used in the combination of ACE inhibitor and aldosterone receptor antagonist. Such diuretic agent may be selected from several known classes, such as thiazides and related sulfonamides, potassium-sparing diuretics, loop diuretics and organic mercurial diuretics.



Angiotensin II receptor antagonist compounds suitable for use in the combination therapy are described in Table II, below. Preferred compounds for use in the combination therapy may be generally characterized structurally as having two portions. A first portion constitutes a mono-aryl-alkyl moiety, or a bi-aryl-alkyl moiety, or a mono-heteroaryl-alkyl moiety, or a bi-heteroaryl-alkyl moiety. A second portion constitutes a heterocyclic moiety or an open chain hetero-atom-containing moiety.

Typically, the first-portion mono/bi-aryl/heteroaryl-alkyl moiety is attached to the second portion heterocyclic/open-chain moiety through the alkyl group of the mono/bi-aryl/heteroaryl-alkyl moiety to any substitutable position on the heterocyclic/open-chain moiety second portion. Suitable first-portion mono/bi-aryl/heteroaryl-alkyl moieties are defined by any of the various moieties listed under Formula I:



wherein the abbreviated notation used in the moieties of Formula I is defined as follows:

"Ar" means a five or six-membered carbocyclic ring system consisting of one ring or two fused rings, with such ring or rings being typically fully unsaturated but which also may be partially or fully saturated. "Phenyl" radical most typically exemplifies "Ar".

"Het" means a monocyclic or bicyclic fused ring

system having from five to eleven ring members, and having at least one of such ring members being a hetero atom selected from oxygen, nitrogen and sulfur, and with such ring system containing up to six of such hetero atoms as ring members.

"Alk" means an alkyl radical or alkylene chain, linear or branched, containing from one to about five carbon atoms. Typically, "Alk" means "methylene", i.e., -CH<sub>2</sub>-.



10

"L" designates a single bond or a bivalent linker moiety selected from carbon, oxygen and sulfur. When "L" is carbon, such carbon has two hydrido atoms attached thereto.

15

Suitable second-portion heterocyclic moieties of the angiotensin II antagonist compounds, for use in the combination therapy, are defined by any of the various moieties listed under Formula IIa or IIb:

17



wherein each of X<sup>1</sup> through X<sup>6</sup> is selected from -CH=, -CH<sub>2</sub>-,  
 5 -N=, -NH-, O, and S, with the proviso that at least one of  
 X<sup>1</sup> through X<sup>6</sup> in each of Formula IIa and Formula IIb must be  
 a hetero atom. The heterocyclic moiety of Formula IIa or  
 IIb may be attached through a bond from any ring member of  
 the Formula IIa or IIb heterocyclic moiety having a  
 10 substitutable or a bond-forming position.

Examples of monocyclic heterocyclic moieties of  
 Formula IIa include thienyl, furyl, pyranyl, pyrrolyl,  
 imidazolyl, triazolyl, pyrazolyl, pyridyl, pyrazinyl,  
 15 pyrimidinyl, pyridazinyl, isothiazolyl, isoxazolyl,  
 furazanyl, pyrrolidinyl, pyrrolinyl, furanyl, thiophenyl,  
 isopyrrolyl, 3-isopyrrolyl, 2-isoimidazolyl, 1,2,3-  
 triazolyl, 1,2,4-triazolyl, 1,2-dithiolyl, 1,3-dithiolyl,  
 1,2,3-oxathiolyl, oxazolyl, thiazolyl, 1,2,3-oxadiazolyl,  
 20 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,  
 1,2,3,4-oxatriazolyl, 1,2,3,5-oxatriazolyl, 1,2,3-  
 dioxazolyl, 1,2,4-dioxazolyl, 1,3,2-dioxazolyl, 1,3,4-  
 dioxazolyl, 1,2,5-oxathiazolyl, 1,3-oxathiolyl, 1,2-pyranyl,  
 1,4-pyranyl, 1,2-pyronyl, 1,4-pyronyl, pyridinyl,  
 25 piperazinyl, s-triazinyl, as-triazinyl, v-triazinyl, 1,2,4-  
 oxazinyl, 1,3,2-oxazinyl, 1,3,6-oxazinyl, 1,2,6-oxazinyl,  
 1,4-oxazinyl, *q*-isoxazinyl, *p*-isoxazinyl, 1,2,5-  
 oxathiazinyl, 1,2,6-oxathiazinyl, 1,4,2-oxadiazinyl,

1,3,5,2-oxadiazinyl, morpholinyl, azepinyl, oxepinyl, thiepinyl and 1,2,4-diazepinyl.

Examples of bicyclic heterocyclic moieties of

- 5 Formula IIb include benzo[b]thienyl, isobenzofuranyl, chromenyl, indoliziny, isoindolyl, indolyl, indazolyl, purinyl, quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnoliny, pteridinyl, isochromanyl, chromanyl, thieno[2,3-b]furanyl, 10 2H-furo[3,2-b]pyranyl, 5H-pyrido[2,3-d][1,2]oxazinyl, 1H-pyrazolo[4,3-d]oxazolyl, 4H-imidazo[4,5-d]thiazolyl, pyrazino[2,3-d]pyridazinyl, imidazo[2,1-b]thiazolyl, cyclopenta[b]pyranyl, 4H-[1,3]oxathiole-[5,4-b]pyrrolyl, thieno[2,3-b]furanyl, imidazo[1,2-b][1,2,4]triazinyl and 15 4H-1,3-dioxolo[4,5-d]imidazolyl.

The angiotensin II receptor antagonist compounds, as provided by the first-and-second-portion moieties of Formula I and II, are further characterized by an acidic moiety attached to either of said first-and-second-portion moieties. Preferably this acidic moiety is attached to the first-portion moiety of Formula I and is defined by Formula III:

25  $-U_nA$  (III)

wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein 30 U is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms.


35

The phrase "acidic group selected to contain at least one acidic hydrogen atom", as used to define the  $-U_nA$

moiety, is intended to embrace chemical groups which, when attached to any substitutable position of the Formula I-IIa/b moiety, confers acidic character to the compound of Formula I-IIa/b. "Acidic character" means proton-donor capability, that is, the capacity of the compound of Formula I-IIa/b to be a proton donor in the presence of a proton-receiving substance such as water. Typically, the acidic group should be selected to have proton-donor capability such that the product compound of Formula I-IIa/b has a  $pK_a$  in a range from about one to about twelve. More typically, the Formula I-IIa/b compound would have a  $pK_a$  in a range from about two to about seven. An example of an acidic group containing at least one acidic hydrogen atom is carboxyl group (-COOH). Where n is zero and A is -COOH, in the  $-U_nA$  moiety, such carboxyl group would be attached directly to one of the Formula I-IIa/b positions. The Formula I-IIa/b compound may have one  $-U_nA$  moiety attached at one of the Formula I-IIa/b positions, or may have a plurality of such  $-U_nA$  moieties attached at more than one of the Formula I-IIa/b positions. There are many examples of acidic groups other than carboxyl group, selectable to contain at least one acidic hydrogen atom. Such other acidic groups may be collectively referred to as "bioisosteres of carboxylic acid" or referred to as "acidic bioisosteres". Specific examples of such acidic bioisosteres are described hereinafter. Compounds of Formula I-IIa/b may have one or more acidic protons and, therefore, may have one or more  $pK_a$  values. It is preferred, however, that at least one of these  $pK_a$  values of the Formula I-IIa/b compound as conferred by the  $-U_nA$  moiety be in a range from about two to about seven. The  $-U_nA$  moiety may be attached to one of the Formula I-IIa/b positions through any portion of the  $-U_nA$  moiety which results in a Formula I-IIa/b compound being relatively stable and also having a labile or acidic proton to meet the foregoing  $pK_a$  criteria. For example, where the  $-U_nA$  acid moiety is tetrazole, the tetrazole is typically attached at


the tetrazole ring carbon atom.

For any of the moieties embraced by Formula I and Formula II, such moieties may be substituted at any  
 5 substitutable position by one or more radicals selected from hydrido, hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, haloalkyl, halo, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano,  
 10 cyanoamino, nitro, alkylcarbonyloxy, alkoxycarbonyloxy, alkylcarbonyl, alkoxycarbonyl, aralkoxycarbonyl, carboxyl, mercapto, mercaptocarbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl,  
 15 arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino and amido radicals of the formula



20

wherein W is oxygen atom or sulfur atom; wherein each of R<sup>1</sup> through R<sup>5</sup> is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, YR<sup>6</sup> and




25

wherein Y is selected from oxygen atom and sulfur atom and R<sup>6</sup> is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R<sup>1</sup>, R<sup>2</sup>,  
 30 R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup> and R<sup>8</sup> is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl,

arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup> and R<sup>8</sup> is further independently selected from amino and amido radicals of the formula

5



wherein W is oxygen atom or sulfur atom;

wherein each of R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is

- 10 independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R<sup>2</sup> and R<sup>3</sup> taken together and each of R<sup>4</sup> and R<sup>5</sup> taken together may form a heterocyclic group having five
- 15 to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially
- 20 unsaturated; wherein each of R<sup>2</sup> and R<sup>3</sup> taken together and each of R<sup>7</sup> and R<sup>8</sup> taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more
- 25 hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

- The combination therapy of the invention would be
- 30 useful in treating a variety of circulatory disorders, including cardiovascular disorders, such as hypertension, congestive heart failure, myocardial fibrosis and cardiac hypertrophy. The combination therapy would also be useful with adjunctive therapies. For example, the combination

therapy may be used in combination with other drugs, such as a diuretic, to aid in treatment of hypertension.

Table II, below, contains description of  
5 angiotensin II antagonist compounds which may be used in the  
combination therapy. Associated with each compound listed  
in Table II is a published patent document describing the  
chemical preparation of the angiotensin II antagonist  
compound as well as the biological properties of such  
10 compound. The content of each of these patent documents is  
incorporated herein by reference.




TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

1

WO #91/17148  
pub. 14 Nov 91

2

WO #91/17148  
pub. 14 Nov 91

3


WO #91/17148  
pub. 14 Nov 91


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

4

WO #91/17148  
pub. 14 Nov 91

5

WO #91/17148  
pub. 14 Nov 91

6


WO #91/17148  
pub. 14 Nov 91


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

7.

WO #91/17148  
pub. 14 Nov 91

8

WO #91/17148  
pub. 14 Nov 91

9


WO #91/17148  
pub. 14 Nov 91

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
10		WO #91/17148 pub. 14 Nov 91
11		WO #91/17148 pub. 14 Nov 91
12		WO #91/17148 pub. 14 Nov 91

TABLE II: Angiotensin II Antagonists







Compound #	Structure	Source
13		WO #91/17148 pub. 14 Nov 91
14		WO #91/17148 pub. 14 Nov 91
15		WO #91/17148 pub. 14 Nov 91

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
16		WO #91/17148 pub. 14 Nov 91
17		WO #91/17148 pub. 14 Nov 91
18		WO #91/17148 pub. 14 Nov 91

29

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
19		WO #91/17148 pub. 14 Nov 91
20		WO #91/17148 pub. 14 Nov 91
21		WO #91/17148 pub. 14 Nov 91


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


22

WO #91/17148  
pub. 14 Nov 91

23

WO #91/17148  
pub. 14 Nov 91

24

WO #91/17148  
pub. 14 Nov 91



31


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

25

WO #91/17148  
pub. 14 Nov 91

26

WO #91/17148  
pub. 14 Nov 91

27





WO #91/17148  
pub. 14 Nov 91

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
28		WO #91/17148 pub. 14 Nov 91
29		WO #91/17148 pub. 14 Nov 91
30		WO #91/17148 pub. 14 Nov 91

33


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


31

WO #91/17148  
pub. 14 Nov 91

32

WO #91/17148  
pub. 14 Nov 91

33

WO #91/17148  
pub. 14 Nov 91

34


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


34

WO #91/17148  
pub. 14 Nov 91

35

WO #91/17148  
pub. 14 Nov 91

36

WO #91/17148  
pub. 14 Nov 91

35


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


37

WO #91/17148  
pub. 14 Nov 91




38

WO #91/17148  
pub. 14 Nov 91

39




WO #91/17148  
pub. 14 Nov 91

**TABLE II: Angiotensin II Antagonists**

Compound #	Structure	Source
40		WO #91/17148 pub. 14 Nov 91
41		WO #91/17148 pub. 14 Nov 91
42		WO #91/17148 pub. 14 Nov 91

37

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
43		WO #91/17148 pub. 14 Nov 91
44		WO #91/17148 pub. 14 Nov 91
45		WO #91/17148 pub. 14 Nov 91

SUBSTITUTE SHEET (RULE 26)

38

TABLE II: Angiotensin II Antagonists







Compound #	Structure	Source
46		WO #91/17148 pub. 14 Nov 91
47		WO #91/17148 pub. 14 Nov 91
48		WO #91/17148 pub. 14 Nov 91



TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
49		WO #91/17148 pub. 14 Nov 91
50		WO #91/17148 pub. 14 Nov 91
51		WO #91/17148 pub. 14 Nov 91

40


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


52

WO #91/17148  
pub. 14 Nov 91

53

WO #91/17148  
pub. 14 Nov 91

54

WO #91/17148  
pub. 14 Nov 91

41


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

55

WO #91/17148  
pub. 14 Nov 91

56

WO #91/17148  
pub. 14 Nov 91

57


WO #91/17148  
pub. 14 Nov 91

TABLE II: Angiotensin II Antagonists





Compound #	Structure	Source
58		WO #91/17148 pub. 14 Nov 91
59		WO #91/17148 pub. 14 Nov 91
60		WO #91/17148 pub. 14 Nov 91


TABLE II: Angiotensin II Antagonists.

Compound #	Structure	Source
------------	-----------	--------

61

WO #91/17148  
pub. 14 Nov 91

62

WO #91/17148  
pub. 14 Nov 91

63


WO #91/17148  
pub. 14 Nov 91


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


64

WO #91/17148  
pub. 14 Nov 91

65

WO #91/17148  
pub. 14 Nov 91

66

WO #91/17148  
pub. 14 Nov 91

45


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

67

WO #91/17148  
pub. 14 Nov 91

68

WO #91/17148  
pub. 14 Nov 91

69


WO #91/17148  
pub. 14 Nov 91


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


70

WO #91/17148  
pub. 14 Nov 91

71

WO #91/17148  
pub. 14 Nov 91

72

WO #91/17148  
pub. 14 Nov 91



47


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


73

WO #91/17148  
pub. 14 Nov 91

74

WO #91/17148  
pub. 14 Nov 91

75

WO #91/17148  
pub. 14 Nov 91

48

TABLE II: Angiotensin II Antagonists



Compound #	Structure	Source
76		WO #91/17148 pub. 14 Nov 91
77		WO #91/17148 pub. 14 Nov 91

TABLE II: Angiotensin II Antagonists










Compound #	Structure	Source
78		WO #91/18888 pub.
79		WO #91/18888 pub.
80		WO #91/18888 pub.

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
81		WO #91/18888 pub.
82		WO #91/18888 pub.
83		WO #91/18888 pub.

51

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
84		WO #91/18888 pub.
85		WO #91/18888 pub.
86		WO #91/18888 pub.

52

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
87		WO #91/18888 pub.
88		WO #91/18888 pub.
89		WO #91/18888 pub.

TABLE II: Angiotensin II Antagonists







Compound #	Structure	Source
90		WO #91/18888 pub.
91		WO #91/18888 pub.
92		WO #91/18888 pub.

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
93		WO #91/18888 pub.
94		WO #91/18888 pub.
95		WO #91/18888 pub.



55

TABLE II: Angiotensin II Antagonists







Compound #	Structure	Source
96		WO #91/18888 pub.
97		WO #91/18888 pub.
98		WO #91/18888 pub.






TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
102		WO #91/18888 pub.
103		WO #91/18888 pub.
104		WO #91/18888 pub.

58

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
105		WO #91/18888 pub.
106		WO #91/18888 pub.
107		WO #91/18888 pub.

59

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
108		WO #91/19715 pub. 26 Dec 91
109		WO #91/19715 pub. 26 Dec 91
110		WO #91/19715 pub. 26 Dec 91

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
111		WO #91/19715 pub. 26 Dec 91
112		WO #91/19715 pub. 26 Dec 91
113		WO #91/19715 pub. 26 Dec 91

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
114		WO #91/19715 pub. 26 Dec 91
115		WO #91/19715 pub. 26 Dec 91
116		WO #91/19715 pub. 26 Dec 91


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

117

WO #91/19715  
pub. 26 Dec 91

118

WO #91/19715  
pub. 26 Dec 91

119


WO #91/19715  
pub. 26 Dec 91




TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

120

WO #91/19715  
pub. 26 Dec 91

121

WO #91/19715  
pub. 26 Dec 91

122


WO #91/19715  
pub. 26 Dec 91


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

123

WO #91/19715  
pub. 26 Dec 91

124

WO #91/19715  
pub. 26 Dec 91

125


WO #91/19715  
pub. 26 Dec 91

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
126		WO #92/05161 pub. 2 Apr 92
127		WO #92/05161 pub. 2 Apr 92
128		WO #92/05161 pub. 2 Apr 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

129

WO #92/05161  
pub. 2 Apr 92

130

WO #92/05161  
pub. 2 Apr 92

131


WO #92/05161  
pub. 2 Apr 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
132		WO #92/07834 pub. 14 May 92
133		WO #92/07834 pub. 14 May 92
134		WO #92/07834 pub. 14 May 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
135		WO #92/07834 pub. 14 May 92
136		WO #92/07834 pub. 14 May 92
137		WO #92/07834 pub. 14 May 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
138		WO #92/07834 pub. 14 May 92
139		WO #92/11255 pub. 9 Jul 92
140		WO #92/11255 pub. 9 Jul 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
141		WO #92/11255 pub. 9 Jul 92
142		WO #92/11255 pub. 9 Jul 92
143		WO #92/11255 pub. 9 Jul 92





TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
147		WO #92/15577 pub. 17 Sep 92
148		WO #92/15577 pub. 17 Sep 92
149		WO #92/15577 pub. 17 Sep 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
150		WO #92/16523 pub. 1 Oct 92
151		WO #92/16523 pub. 1 Oct 92
152		WO #92/16523 pub. 1 Oct 92

TABLE II: Angiotensin II Antagonists





Compound #	Structure	Source
153		WO #92/16523 pub. 1 Oct 92
154		WO #92/16523 pub. 1 Oct 92
155		WO #92/16523 pub. 1 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------


156

WO #92/16523  
pub. 1 Oct 92

157

WO #92/16523  
pub. 1 Oct 92

158

WO #92/16523  
pub. 1 Oct 92

76


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

159

WO #92/16523  
pub. 1 Oct 92

160

WO #92/16523  
pub. 1 Oct 92

161


WO #92/16523  
pub. 1 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

162

WO #92/16523  
pub. 1 Oct 92

163

WO #92/16523  
pub. 1 Oct 92

164


WO #92/16523  
pub. 1 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

165

WO #92/16523  
pub. 1 Oct 92

166

WO #92/16523  
pub. 1 Oct 92

167


WO #92/16523  
pub. 1 Oct 92



TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
168		WO #92/16523 pub. 1 Oct 92
169		WO #92/16523 pub. 1 Oct 92
170		WO #92/16523 pub. 1 Oct 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
171		WO #92/16523 pub. 1 Oct 92
172		WO #92/16523 pub. 1 Oct 92
173		WO #92/16523 pub. 1 Oct 92

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
174	 <chem>COCC1=CN=C(NCCC)N1CCc2ccncc2-c3ccccc3C4=NN=CN4</chem>	WO #92/16523 pub. 1 Oct 92
175	 <chem>COCC1=CN=C(NCCC)N1CCc2ccncc2-c3ccccc3C4=NN=CN4</chem>	WO #92/16523 pub. 1 Oct 92
176	 <chem>CCOC1=CN=C(NCCC)N1CCc2ccncc2-c3ccccc3C4=NN=CN4</chem>	WO #92/16523 pub. 1 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

177

WO #92/16523  
pub. 1 Oct 92

178

WO #92/16523  
pub. 1 Oct 92

179


WO #92/16523  
pub. 1 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

180

WO #92/16523  
pub. 1 Oct 92

181

WO #92/16523  
pub. 1 Oct 92

182





WO #92/16523  
pub. 1 Oct 92

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
183		WO #92/16523 pub. 1 Oct 92
184		WO #92/16523 pub. 1 Oct 92
185		WO #92/17469 pub. 15 Oct 92

85

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
186		WO #92/17469 pub. 15 Oct 92
187		WO #92/17469 pub. 15 Oct 92
188		WO #92/17469 pub. 15 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

189

WO #92/17469  
pub. 15 Oct 92

190

WO #92/17469  
pub. 15 Oct 92

191


WO #92/17469  
pub. 15 Oct 92



TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
192		WO #92/17469 pub. 15 Oct 92
193		WO #92/17469 pub. 15 Oct 92
194		WO #92/17469 pub. 15 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

195

WO #92/17469  
pub. 15 Oct 92

196

WO #92/17469  
pub. 15 Oct 92

197


WO #92/17469  
pub. 15 Oct 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
198		WO #92/17469 pub. 15 Oct 92
199		WO #92/17469 pub. 15 Oct 92
200		WO #92/17469 pub. 15 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

201

WO #92/17469  
pub. 15 Oct 92

202

WO #92/17469  
pub. 15 Oct 92

203


WO #92/17469  
pub. 15 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


204

WO #92/17469  
pub. 15 Oct 92

205

WO #92/17469  
pub. 15 Oct 92

206

WO #92/17469  
pub. 15 Oct 92

92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

207

WO #92/17469  
pub. 15 Oct 92

208

WO #92/17469  
pub. 15 Oct 92

209


WO #92/17469  
pub. 15 Oct 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
210		WO #92/17469 pub. 15 Oct 92
211		WO #92/17469 pub. 15 Oct 92
212		WO #92/17469 pub. 15 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

213

WO #92/17469  
pub. 15 Oct 92

214

WO #92/17469  
pub. 15 Oct 92

215


WO #92/17469  
pub. 15 Oct 92




TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

216

WO #92/17469  
pub. 15 Oct 92

217

WO #92/17469  
pub. 15 Oct 92

218


WO #92/17469  
pub. 15 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

219

WO #92/17469  
pub. 15 Oct 92

220

WO #92/17469  
pub. 15 Oct 92

221


WO #92/17469  
pub. 15 Oct 92


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

222

WO #92/17469  
pub. 15 Oct 92

223

WO #92/17469  
pub. 15 Oct 92

224


WO #92/17469  
pub. 15 Oct 92

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
225	 <chem>CCCCN1C(=O)C=C(N1Cc2ccc(cc2)-c3ccncc3C4=NN=N4)C5=CC=CC=C5</chem>	WO #92/17469 pub. 15 Oct 92
226	 <chem>CCCCN1C(=O)C=C(N1Cc2ccc(cc2)-c3ccncc3C4=NN=N4)C5=CC=CC=C5</chem>	WO #92/17469 pub. 15 Oct 92
227	 <chem>CCCCN1C(=O)C=C(N1Cc2ccc(cc2)-c3ccncc3C4=NN=N4)C5=CC=CC=C5</chem>	WO #92/17469 pub. 15 Oct 92

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------

228		
-----	---	--

229		
-----	--	--


230		
-----	---	--

100


TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------


231



232



233



101

TABLE II: Angiotensin II Antagonists


Compound #	Structure	Source
234		
235		
236		

**SUBSTITUTE SHEET (RULE 26)**


TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------

237



238



239





WO #92/18092  
pub. 29 Oct 92



TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
240		WO #92/18092 pub. 29 Oct 92
241		WO #92/18092 pub. 29 Oct 92
242		WO #92/18092 pub. 29 Oct 92

104


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


243

WO #92/18092  
pub. 29 Oct 92

244




WO #92/18092  
pub. 29 Oct 92

245

WO #92/18092  
pub. 29 Oct 92

105

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
246		WO #92/18092 pub. 29 Oct 92
247		WO #92/18092 pub. 29 Oct 92
248		WO #92/18092 pub. 29 Oct 92

106

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
249		WO #92/18092 pub. 29 Oct 92
250		WO #92/18092 pub. 29 Oct 92
251		WO #92/18092 pub. 29 Oct 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
252		WO #92/18092 pub. 29 Oct 92
253		WO #92/18092 pub. 29 Oct 92
254		WO #92/18092 pub. 29 Oct 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
255		WO #92/18092 pub. 29 Oct 92
256		WO #92/18092 pub. 29 Oct 92
257		WO #92/18092 pub. 29 Oct 92

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
258		WO #92/18092 pub. 29 Oct 92
259		WO #92/18092 pub. 29 Oct 92
260		WO #92/18092 pub. 29 Oct 92

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
261	 <chem>CCCC1=CN=C(C2=CC=C(C=C2)C)N1CC3=CC=C(C=C3C4=CC=CC=C4C5=NN=CN5)N</chem>	WO #92/18092 pub. 29 Oct 92
262	 <chem>CCCC1=CN=C(C2=CC(=C)C=C2C)N1CC3=CC=C(C=C3C4=CC=CC=C4C5=NN=CN5)N</chem>	WO #92/18092 pub. 29 Oct 92
263	 <chem>CCCC1=CN=C(C2=CC=C(C=C2)CN(C)C)N1CC3=CC=C(C=C3C4=CC=CC=C4C5=NN=CN5)N</chem>	WO #92/18092 pub. 29 Oct 92






111

**TABLE II: Angiotensin II Antagonists**

Compound #	Structure	Source
264		WO #92/18092 pub. 29 Oct 92
265		WO #92/18092 pub. 29 Oct 92
266		WO #92/18092 pub. 29 Oct 92

112

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
267		WO #92/18092 pub. 29 Oct 92
268		WO #92/18092 pub. 29 Oct 92
269		WO #92/18092 pub. 29 Oct 92

113


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


270

WO #92/18092  
pub. 29 Oct 92

271




PCT/US94/02156  
filed 8 Mar 94

272

PCT/US94/02156  
filed 8 Mar 94

114

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
273		PCT/US94/02156 filed 8 Mar 94
274		PCT/US94/02156 filed 8 Mar 94
275		PCT/US94/02156 filed 8 Mar 94

SUBSTITUTE SHEET (RULE 26)

115


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


276

PCT/US94/02156  
filed 8 Mar 94

277

PCT/US94/02156  
filed 8 Mar 94

278

PCT/US94/02156  
filed 8 Mar 94

116


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

279

PCT/US94/02156  
filed 8 Mar 94

280

WO #91/17148  
pub. 14 Nov. 91

117

TABLE II: Angiotensin II Antagonists





Compound #	Structure	Source
281		EP #475,206 pub. 18 Mar 92
282		WO #93/18035 pub. 16 Sep 93
283		WO #93/17628 pub. 16 Sep 93
284		WO #93/17681 pub. 16 Sep 93


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

285

EP #513,533  
pub. 19 Nov 92

286

EP #535,463  
pub. 07 Apr 93

287





EP #535,465  
pub. 07 Apr 93



TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
288		EP #539,713 pub. 05 May 93
289		EP #542,059 pub. 19 May 93
290		EP #05 557,843 pub. 01 Sep 93




120

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
291	 <chem>CCOC(=O)c1ccc(cc1)CN2C(=NC(C2)C3=CC=C(C=C3)COC)CC4</chem>	EP #563,705 pub. 06 Oct 93
292	 <chem>CCCC1=C(C(=O)O)N(C1)CN2C(=CC=C2)C(=O)O</chem>	EP #562,261 pub. 26 Sep 93
293	 <chem>CCOC(=O)c1ccc(cc1)CN2C(=NC(C2)C3=CC=C(C=C3)COC)CC4</chem>	EP #05 557,843 pub. 15 Sep 93

121

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
294		EP #560,163 pub. 15 Sep 93
295		EP #564,788 pub. 13 Oct 93
296		EP #565,986 pub. 20 Oct 93

SUBSTITUTE SHEET (RULE 26)

122

TABLE II: Angiotensin II Antagonists







Compound #	Structure	Source
297		EP #0,569,795 pub. 18 Nov 93
298		EP #0,569,794 pub. 18 Nov 93
299		EP #0,578,002 pub. 12 Jan 94

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
300	 <chem>CCCC1=C(Cl)N(C1C(=O)O)CC2=CC=C(C=C2C3CCCCC3C(=O)NS(=O)(=O)C4=CC=C(C=C4)C)C3CCCCC3C(=O)NS(=O)(=O)C4=CC=C(C=C4)C</chem>	EP #581,003 pub. 02 Feb 94
301	 <chem>CCCC1=C(Cl)N(C1C(=O)O)CC2=CC=C(C=C2C3CCCCC3C(=O)NS(=O)(=O)C4=CC=C(C=C4)C)C3CCCCC3C(=O)NS(=O)(=O)C4=CC=C(C=C4)C</chem>	EP #392,317 pub. 17 Oct 90
302	 <chem>CCCC1=C(Cl)N(C1C(=O)O)CC2=CC=C(C=C2C3CCCCC3C(=O)NS(=O)(=O)C4=CC=C(C=C4)C)C3CCCCC3C(=O)NS(=O)(=O)C4=CC=C(C=C4)C</chem>	EP #392,317 pub. 17 Oct 90





124

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
303		EP #502,314 pub. 09 Sep 92
304		EP #468,740 pub. 29 Jan 92
305		EP #470,543 pub. 12 Feb 92

125

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
306		EP #502,314 pub. 09 Sep 92
307		EP #529,253 pub. 03 Mar 93
308		EP #543,263 pub. 26 May 93
309		EP #552,765 pub. 28 Jul 93

126

TABLE II: Angiotensin II Antagonists







Compound #	Structure	Source
310		EP #558,825 pub. 18 Aug 93
311		EP #556,789 pub. 25 Aug 93
312		EP #560,330 pub. 15 Sep 93







TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
313		EP #566,020 pub. 20 Oct 93
314		EP #581,166 pub. 02 Feb 94
315		WO #94/01436 pub. 20 Jan 94

128

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
316		EP #253,310 pub. 20 Jan 88
317		EP #324,377 pub. 19 Jul 89
318		US #5,043,349 issued 27 Aug 91
319		WO #91/00281 pub. 10 Jan 91

129


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


320

US #5,015,651  
pub. 14 May 91


321



322




WO #92/00977  
pub. 23 Jan 92

323






130

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
324		WO #93/04046 pub. 04 Mar 93
325		WO #93/10106 pub. 27 May 93
326		US #5,219,856 pub. 15 Jun 93

131

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
327		US #5,260,325 pub. 09 Nov 93
328		US #5,264,581 pub. 23 Nov 93
329		EP #400,974 pub. 5 Dec 90





132

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
330	 <chem>CC1=CC=C(C=C1)N2C(=O)N(C2C3=CC=C(C=C3)C4=CC=CC=C4C5=CC=CC=C5C6=NN=CN6)C7=CC=C(C=C7)C</chem>	EP #411,766 pub. 06 Feb 91
331	 <chem>Clc1ccccc1N2C(=O)N(C2C3=CC=C(C=C3)C4=CC=CC=C4C5=CC=CC=C5C6=NN=CN6)C7=CC=C(C=C7)C</chem>	EP #412,594 pub. 13 Feb 91
332	 <chem>CC1=CC=C(C=C1)N2C(=O)N(C2C3=CC=C(C=C3)C4=CC=CC=C4C5=CC=CC=C5C6=NN=CN6)C7=CC=C(C=C7)C</chem>	EP #419,048 pub. 27 Mar 91




133

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
333		WO #91/12,001 pub. 22 Aug 91
334		WO #91/11,999 pub. 22 Aug 91
335		WO #91/11,909 pub. 22 Aug 91
336		WO #91/12,002 pub. 22 Aug 91

134

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
337		US 5,053,329 pub. 01 Oct 91
338		US #5,057,522 pub. 15 Oct 91
339		WO #91/15,479 pub. 17 Oct 91




135

TABLE II: Angiotensin II Antagonists


Compound #	Structure	Source
------------	-----------	--------

340




EP #456,510  
pub. 13 Nov 91

341



EP #467,715  
pub. 22 Jan 92

342



US #5,087,702  
pub. 11 Feb 92

136


TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source

343

EP #479,479  
pub. 08 Apr 92

344



345








EP #481,614  
pub. 22 Apr 92

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
346		EP #490,587 pub. 17 Jun 92
347		US #5,128,327 pub. 07 Jul 92
348		US #5,132,216 pub. 21 Jul 92

138

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
349		EP #497,516 pub. 05 Aug 92
350		EP #502,725 pub. 09 Sep 92
351		EP #502,575 pub. 09 Sep 92

139

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
352	 <chem>CCCCNC(=O)Cc1ccc(cc1)-c2ccccc2C3=CN=C(C4=CC=CC=C4C5=CC=CC=C5C(=O)NCCC)N3</chem>	EP #503,838 pub. 16 Sep 92
353	 <chem>CCCCNC(=O)Cc1ccc(cc1)-c2ccccc2C3=NN=C(C4=CC=CC=C4C5=CC=CC=C5C(=O)NCCC)N3</chem>	EP #505,111 pub. 23 Sep 92
354	 <chem>CCCCNC(=O)Cc1ccc(cc1)-c2ccccc2C3=NN=C(C4=CC=CC=C4C5=CC=CC=C5C(=O)NCCC)N3</chem>	EP #505,098 pub. 23 Sep 92

140

TABLE II: Angiotensin II Antagonists







Compound #	Structure	Source
355		EP #597,594 pub. 07 Oct 92
356		EP #508,723 pub. 14 Oct 92
357		

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
358		EP #512,675 pub. 11 Nov 92
359		EP #512,676 pub. 11 Nov 92
360		EP #512,870 pub. 11 Nov 92

142

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
361		EP #513,979 pub. 19 Nov 92
362		WO #92/20,660 pub. 26 Nov 92
363		WO #92/20,661 pub. 26 Nov 92




TABLE II: Angiotensin II Antagonists

Compound #


Structure

Source


364

WO 92/20,662  
pub. 26 Nov 92

365

WO #92/20,687  
pub. 26 Nov 92




366

EP #517,357  
pub. 09 Dec 92






145

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
370		US #5,202,322 pub. 13 Apr 93
371		EP #537,937 pub. 21 Apr 93
372		US #5,217,882 pub. 08 Jun 93

**TABLE II: Angiotensin II Antagonists**

Compound #	Structure	Source
373		US #5,214,153 pub. 25 May 93
374		US #5,218,125 pub. 08 Jun 93
375		US #5,236,928 pub. 17 Aug 93

147




TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
376		US #5,240,938 pub. 31 Aug 93
377		GB #2,264,709 pub. 08 Sep 93
378		GB #2,264,710 pub. 08 Sep 93

148

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
------------	-----------	--------



149

TABLE II: Angiotensin II Antagonists




Compound #	Structure	Source
------------	-----------	--------



150

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------








151




TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------







152

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
390		EP #425,211 pub. 02 May 91
391		EP #427,463 pub. 15 May 91
392		WO #92/00068 pub. 09 Jan 92




**TABLE II: Angiotensin II Antagonists**

Compound #	Structure	Source
393		WO #92/02,510 pub. 20 Feb 92
394		WO #92/09278 pub. 11 Jun 92
395		WO #92/10181 pub. 25 Jun 92
396		

154

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------




155


TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------


400



401



402




156


TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------


403

WO #92/10097  
pub. 25 Jun 92

404



405




157


TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------

406




407



WO #92/20651  
pub. 26 Nov 92

408



WO #93/03018  
pub. 18 Feb 93


158

TABLE II: Angiotensin II Antagonists


Compound #

Structure


Source



WO #94/00120  
pub. 06 Jan 94



EP #459,136  
pub. 04 Dec 91






EP #411,507  
pub. 05 Feb 91






159

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
412		EP #425,921 pub. 08 May 91
413		EP #430,300 pub. 05 Jun 91
414		EP #434,038 pub. 26 Jun 91




160

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
415		EP #442,473 pub. 21 Aug 91
416		EP #443,568 pub. 28 Aug 91
417		EP #459,136 pub. 04 Dec 91




161

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
418		EP #483,683 pub. 05 May 92
419		EP #518,033 pub. 16 Dec 92
420		EP #520,423 pub. 30 Dec 92




162

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
421		EP #546,358 pub. 16 Jun 93
422		WO #93/00341 pub. 07 Jan 93
423		WO #92/06081 pub. 16 Apr 92

163




TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
424		WO #93/00341 pub. 07 Jan 93
425		US #5,210,204 pub. 11 May 93
426		EP #343,654 pub. 29 Nov 89

164

TABLE II: Angiotensin II Antagonists

Compound #	Structure	Source
------------	-----------	--------



The term "hydrido" denotes a single hydrogen atom (H). This hydrido group may be attached, for example, to an oxygen atom to form a hydroxyl group; or, as another example, one hydrido group may be attached to a carbon atom

5 to form a  $\begin{array}{c} \diagup \\ \text{CH}- \\ \diagdown \end{array}$  group; or, as another example, two hydrido atoms may be attached to a carbon atom to form a -CH<sub>2</sub>- group. Where the term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "hydroxyalkyl", the term "alkyl" embraces linear or branched  
10 radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about five carbon atoms. The  
15 term "cycloalkyl" embraces cyclic radicals having three to about ten ring carbon atoms, preferably three to about six carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is  
20 substituted with one or more halo groups, preferably selected from bromo, chloro and fluoro. Specifically embraced by the term "haloalkyl" are monohaloalkyl, dihaloalkyl and polyhaloalkyl groups. A monohaloalkyl group, for example, may have either a bromo, a chloro, or a  
25 fluoro atom within the group. Dihalalkyl and polyhaloalkyl groups may be substituted with two or more of the same halo groups, or may have a combination of different halo groups. A dihaloalkyl group, for example, may have two fluoro atoms, such as difluoromethyl and difluorobutyl groups, or two  
30 chloro atoms, such as a dichloromethyl group, or one fluoro atom and one chloro atom, such as a fluoro-chloromethyl group. Examples of a polyhaloalkyl are trifluoromethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, perfluoroethyl and 2,2,3,3-tetrafluoropropyl groups. The term "difluoroalkyl"  
35 embraces alkyl groups having two fluoro atoms substituted on any one or two of the alkyl group carbon atoms. The terms "alkylol" and "hydroxyalkyl" embrace linear or branched

alkyl groups having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl groups. The term "alkenyl" embraces linear or branched radicals having two to about twenty carbon atoms, preferably three to about ten carbon atoms, and containing at least one carbon-carbon double bond, which carbon-carbon double bond may have either cis or trans geometry within the alkenyl moiety. The term "alkynyl" embraces linear or branched radicals having two to about twenty carbon atoms, preferably two to about ten carbon atoms, and containing at least one carbon-carbon triple bond. The term "cycloalkenyl" embraces cyclic radicals having three to about ten ring carbon atoms including one or more double bonds involving adjacent ring carbons. The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms, such as methoxy group. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy groups attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl groups. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy or haloalkoxyalkyl groups. The term "alkylthio" embraces radicals containing a linear or branched alkyl group, of one to about ten carbon atoms attached to a divalent sulfur atom, such as a methylthio group. Preferred aryl groups are those consisting of one, two, or three benzene rings. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl and biphenyl. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenyl-ethyl, phenylbutyl and diphenylethyl. The terms "benzyl" and "phenylmethyl" are interchangeable. The terms "phenalkyl" and "phenylalkyl" are interchangeable. An example of "phenalkyl" is "phenethyl" which is interchangeable with "phenylethyl". The terms "alkylaryl", "alkoxyaryl" and "haloaryl" denote, respectively, the substitution of one or more "alkyl",



"alkoxy" and "halo" groups, respectively, substituted on an "aryl" nucleus, such as a phenyl moiety. The terms "aryloxy" and "arylthio" denote radicals respectively, provided by aryl groups having an oxygen or sulfur atom through which the radical is attached to a nucleus, examples of which are phenoxy and phenylthio. The terms "sulfinyl" and "sulfonyl", whether used alone or linked to other terms, denotes, respectively, divalent radicals SO and SO<sub>2</sub>. The term "aralkoxy", alone or within another term, embraces an aryl group attached to an alkoxy group to form, for example, benzyloxy. The term "acyl" whether used alone, or within a term such as acyloxy, denotes a radical provided by the residue after removal of hydroxyl from an organic acid, examples of such radical being acetyl and benzoyl. "Lower alkanoyl" is an example of a more preferred sub-class of acyl. The term "amido" denotes a radical consisting of nitrogen atom attached to a carbonyl group, which radical may be further substituted in the manner described herein. The term "monoalkylaminocarbonyl" is interchangeable with "N-alkylamido". The term "dialkylaminocarbonyl" is interchangeable with "N,N-dialkylamido". The term "alkenylalkyl" denotes a radical having a double-bond unsaturation site between two carbons, and which radical may consist of only two carbons or may be further substituted with alkyl groups which may optionally contain additional double-bond unsaturation. The term "heteroaryl", where not otherwise defined before, embraces aromatic ring systems containing one or two hetero atoms selected from oxygen, nitrogen and sulfur in a ring system having five or six ring members, examples of which are thienyl, furanyl, pyridinyl, thiazolyl, pyrimidyl and isoxazolyl. Such heteroaryl may be attached as a substituent through a carbon atom of the heteroaryl ring system, or may be attached through a carbon atom of a moiety substituted on a heteroaryl ring-member carbon atom, for example, through the methylene substituent of imidazolemethyl moiety. Also, such heteroaryl may be attached through a ring nitrogen atom as long as aromaticity

of the heteroaryl moiety is preserved after attachment. For any of the foregoing defined radicals, preferred radicals are those containing from one to about ten carbon atoms.

5           Specific examples of alkyl groups are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, methylbutyl, dimethylbutyl and neopentyl. Typical alkenyl and alkynyl groups may have one unsaturated bond, such as an allyl group, or may have a  
10           plurality of unsaturated bonds, with such plurality of bonds either adjacent, such as allene-type structures, or in conjugation, or separated by several saturated carbons.

          Also included in the combination of the invention  
15           are the isomeric forms of the above-described angiotensin II receptor compounds and the epoxy-free spirolactone-type aldosterone receptor compounds, including diastereoisomers, regioisomers and the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts"  
20           embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts may be prepared from an  
25           inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic,  
30           carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic,  
35           p-hydroxybenzoic, salicylic, phenylacetic, mandelic, embonic (pamoic), methansulfonic, ethanesulfonic, 2-hydroxyethanesulfonic, pantothenic, benzenesulfonic,

toluenesulfonic, sulfanilic, mesylic, cyclohexylaminosulfonic, stearic, algenic,  $\beta$ -hydroxybutyric, malonic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts include

- 5 metallic salts made from aluminium, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by
- 10 conventional means from the corresponding compound by reacting, for example, the appropriate acid or base with such compound.

### BIOLOGICAL EVALUATION

Human congestive heart failure (CHF) is a complex condition usually initiated by vascular hypertension or a myocardial infarction (MI). In order to determine the probable effectiveness of a combination therapy for CHF, it is important to determine the potency of individual components of the combination therapy. Accordingly, in Assays "A" through "C", the angiotensin II receptor antagonist profiles were determined for many of the compounds described in Table II, herein. In Assays "D" and "E", there are described methods for evaluating a combination therapy of the invention, namely, an angiotensin II receptor antagonist of Table II and an epoxy-free spiro lactone-type aldosterone receptor antagonist. The efficacy of the individual drugs, spironolactone and the angiotensin II receptor blocker, and of these drugs given together at various doses, are evaluated in rodent models of hypertension and CHF using surgical alterations to induce either hypertension or an MI. The methods and results of such assays are described below.

#### Assay A: Angiotensin II Binding Activity

Compounds of the invention were tested for ability to bind to the smooth muscle angiotensin II receptor using a rat uterine membrane preparation. Angiotensin II (AII) was purchased from Peninsula Labs.  $^{125}\text{I}$ -angiotensin II (specific activity of 2200 Ci/mmol) was purchased from Du Pont-New England Nuclear. Other chemicals were obtained from Sigma Chemical Co. This assay was carried out according to the method of Douglas et al [Endocrinology, 106, 120-124 (1980)]. Rat uterine membranes were prepared from fresh tissue. All procedures were carried out at 4°C. Uteri were stripped of fat and homogenized in phosphate-buffered saline at pH 7.4 containing 5 mM EDTA. The homogenate was

171

centrifuged at 1500 x g for 20 min., and the supernatant was recentrifuged at 100,000 x g for 60 min. The pellet was resuspended in buffer consisting of 2 mM EDTA and 50 mM Tris-HCl (pH 7.5) to a final protein concentration of 4 mg/ml. Assay tubes were charged with 0.25 ml of a solution containing 5 mM MgCl<sub>2</sub>, 2 mM EDTA, 0.5% bovine serum albumin, 50 mM Tris-HCl, pH 7.5 and <sup>125</sup>I-AII (approximately 10<sup>5</sup> cpm) in the absence or in the presence of unlabelled ligand. The reaction was initiated by the addition of membrane protein and the mixture was incubated at 25°C for 60 min. The incubation was terminated with ice-cold 50 mM Tris-HCl (pH 7.5) and the mixture was filtered to separate membrane-bound labelled peptide from the free ligand. The incubation tube and filter were washed with ice-cold buffer. Filters were assayed for radioactivity in a Micromedic gamma counter. Nonspecific binding was defined as binding in the presence of 10 µM of unlabelled AII. Specific binding was calculated as total binding minus nonspecific binding. The receptor binding affinity of an AII antagonist compound was indicated by the concentration (IC<sub>50</sub>) of the tested AII antagonist which gives 50% displacement of the total specifically bound <sup>125</sup>I-AII from the angiotensin II AT<sub>1</sub> receptor. Binding data were analyzed by a nonlinear least-squares curve fitting program. Results are reported in Table III.

25

#### Assay B: In Vitro Vascular Smooth Muscle-Response for AII

The compounds of the invention were tested for antagonist activity in rabbit aortic rings. Male New Zealand white rabbits (2-2.5 kg) were sacrificed using an overdose of pentobarbital and exsanguinated via the carotid arteries. The thoracic aorta was removed, cleaned of adherent fat and connective tissue and then cut into 3-mm ring segments. The endothelium was removed from the rings by gently sliding a rolled-up piece of filter paper into the vessel lumen. The rings were then mounted in a water-jacketed tissue bath, maintained at 37°C, between moveable and fixed ends of a

stainless steel wire with the moveable end attached to an FT03 Grass transducer coupled to a Model 7D Grass Polygraph for recording isometric force responses. The bath was filled with 20 ml of oxygenated (95% oxygen/5% carbon dioxide) Krebs solution of the following composition (mM): 130 NaCl, 15 NaHCO<sub>3</sub>, 15 KCl, 1.2 NaH<sub>2</sub>PO<sub>4</sub>, 1.2 MgSO<sub>4</sub>, 2.5 CaCl<sub>2</sub>, and 11.4 glucose. The preparations were equilibrated for one hour before approximately one gram of passive tension was placed on the rings. Angiotensin II concentration-response curves were then recorded ( $3 \times 10^{-10}$  to  $1 \times 10^{-5}$  M). Each concentration of AII was allowed to elicit its maximal contraction, and then AII was washed out repeatedly for 30 minutes before rechallenging with a higher concentration of AII. Aorta rings were exposed to the test antagonist at  $10^{-5}$  M for 5 minutes before challenging with AII. Adjacent segments of the same aorta ring were used for all concentration-response curves in the presence or absence of the test antagonist. The effectiveness of the test compound was expressed in terms of pA<sub>2</sub> values and were calculated according to H.O. Schild [Br. J. Pharmacol. Chemother., 2,189-206 (1947)]. The pA<sub>2</sub> value is the concentration of the antagonist which increases the EC<sub>50</sub> value for AII by a factor of two. Each test antagonist was evaluated in aorta rings from two rabbits. Results are reported in Table III.

25

Assay C: In Vivo Intragastric Pressor Assay Response for All Antagonists

Male Sprague-Dawley rats weighing 225-300 grams were anesthetized with methohexital (30 mg/kg, i.p.) and catheters were implanted into the femoral artery and vein. The catheters were tunneled subcutaneously to exit dorsally, posterior to the head and between the scapulae. The catheters were filled with heparin (1000 units/ml of saline). The rats were returned to their cage and allowed regular rat chow and water ad libitum. After full recovery from surgery (3-4 days), rats were placed in Lucite holders

and the arterial line was connected to a pressure transducer. Arterial pressure was recorded on a Gould polygraph (mmHg). Angiotensin II was administered as a 30 ng/kg bolus via the venous catheter delivered in a 50  $\mu$ l volume with a 0.2 ml saline flush. The pressor response in mm Hg was measured by the difference from pre-injection arterial pressure to the maximum pressure achieved. The AII injection was repeated every 10 minutes until three consecutive injections yielded responses within 4 mmHg of each other. These three responses were then averaged and represented the control response to AII. The test compound was suspended in 0.5% methylcellulose in water and was administered by gavage. The volume administered was 2 ml/kg body weight. The standard dose was 3 mg/kg. Angiotensin II bolus injections were given at 30, 45, 60, 75, 120, 150, and 180 minutes after gavage. The pressor response to AII was measured at each time point. The rats were then returned to their cage for future testing. A minimum of 3 days was allowed between tests. Percent inhibition was calculated for each time point following gavage by the following formula:  $[(\text{Control Response} - \text{Response at time point}) / \text{Control Response}] \times 100$ . Results are shown in Table III.

25 Assay "D": Hypertensive Rat Model

Male rats are made hypertensive by placing a silver clip with an aperture of 240 microns on the left renal artery, leaving the contralateral kidney untouched. Sham controls undergo the same procedure but without attachment of the clip. One week prior to the surgery, animals to be made hypertensive are divided into separate groups and drug treatment is begun. Groups of animals are administered vehicle, AII antagonist alone, spironolactone alone, and combinations of AII antagonist and spironolactone, at various doses, as follow:

AII Antagonist (mg/kg/day)	Spironolactone (mg/kg/day)	Combination of AII Antagonist & Spironolactone	
		(mg/kg/day)	(mg/kg/day)
3	5	3	5
	20	3	20
	50	3	50
	100	3	100
	200	3	200
10	5	10	5
	20	10	20
	50	10	50
	100	10	100
	200	10	200
30	5	30	5
	20	30	20
	50	30	50
	100	30	100
	200	30	200

After 12 to 24 weeks, systolic and diastolic blood pressure, left ventricular end diastolic pressure, left ventricular dP/dt, and heart rate are evaluated. The hearts are removed, weighed, measured and fixed in formalin. Collagen content of heart sections are evaluated using computerized image analysis of picrosirius stained sections. It would be expected that rats treated with a combination therapy of AII antagonist and spironolactone components, as compared to rats treated with either component alone, will show improvements in cardiac performance.

Assay "E": Myocardial Infarction Rat Model:

15

Male rats are anesthetized and the heart is exteriorized following a left sided thoracotomy. The left anterior descending coronary artery is ligated with a suture. The thorax is closed and the animal recovers. Sham



animals have the suture passed through without ligation. One week prior to the surgery, animals to undergo infarction are divided into separate groups and drug treatment is begun. Groups of animals are administered vehicle, AII antagonist alone, spironolactone alone, and combinations of AII antagonist and spironolactone, at various doses, as follow:

AII Antagonist (mg/kg/day)	Spironolactone (mg/kg/day)	Combination of AII Antagonist & Spironolactone	
		(mg/kg/day)	(mg/kg/day)
3	5	3	5
	20	3	20
	50	3	50
	100	3	100
	200	3	200
10	5	10	5
	20	10	20
	50	10	50
	100	10	100
	200	10	200
30	5	30	5
	20	30	20
	50	30	50
	100	30	100
	200	30	200

After six weeks, systolic and diastolic blood pressure, left ventricular end diastolic pressure, left ventricular dP/dt, and heart rate are evaluated. The hearts are removed, weighed, measured and fixed in formalin. Collagen content of heart sections are evaluated using computerized image analysis of picrosirius stained sections. It would be expected that rats treated with a combination therapy of AII antagonist and spironolactone components, as compared to rats treated with either component alone, will show improvements in cardiac performance.

176

TABLE III

In Vivo and In Vitro Angiotensin II  
Activity of Compounds of the Invention

Test Compound Example #	<sup>1</sup> Assay A		<sup>2</sup> Assay B		<sup>3</sup> Assay C	
	IC <sub>50</sub>		pA <sub>2</sub>	Dose	Inhibition	Duration
	(nM)			(mg/kg)	(%)	(min.)
10	1	NT	NT	NT	NT	NT
	2	95	7.37/7.59	10	95	60
				30	98	90-120
	3	5.4	8.70 ± 0.2	10	50	>180
15				30	100	200+
	4	NT	NT	NT	NT	NT
	5	200	7.48/6.91	30	38	20-30
	6	1300	6.55/6.82	100	90	120
20	7	86	8.01/8.05	30	90	130
	8	17,000	NT	NT	NT	NT
	9	700	6.67/6.12	30	80	75
				100	100	130
25	10	4.9	8.19/7.59	3	86	100
				30	100	240
	11	160	6.45/6.77	NT	NT	NT
	12	6.0	8.66/8.59	NT	NT	NT
30	13	17	8.70/8.85	NT	NT	NT
	14	7.2	8.84/8.71	NT	NT	NT
	15	16	8.31/8.30	NT	NT	NT
	16	6.4	8.95/9.24	NT	NT	NT
30	17	4.0	8.64/8.40	NT	NT	NT
	18	970	6.14/6.09	NT	NT	NT
	19	12,000	5.18/5.35	NT	NT	NT

177

Test Compound	Example #	<sup>1</sup> Assay A	<sup>2</sup> Assay B	<sup>3</sup> Assay C		
		IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition	Duration
		(nM)		(mg/kg)	(%)	(min.)
5	20	78,000	5.89/5.99	100	10	45
	21	87	7.71/7.21	NT	NT	NT
	22	460	6.60/6.46	NT	NT	NT
	23	450	6.48/7.15	NT	NT	NT
	24	10	7.56/7.73	NT	NT	NT
10	25	480	6.80/6.73	NT	NT	NT
	26	3.2	9.83/9.66	10	50	>180
	27	180	NT	NT	NT	NT
	28	570	5.57/6.00	NT	NT	NT
	29	160	NT	NT	NT	NT
15	30	22	7.73/7.88	30	50	>180
	31	14	NT	NT	NT	NT
	32	16	7.68/7.29	NT	NT	NT
	33	630	6.73/6.36	NT	NT	NT
	34	640	5.34/5.69	NT	NT	NT
20	35	41	7.25/7.47	NT	NT	NT
	36	1400	5.92/5.68	NT	NT	NT
	37	340	6.90/6.85	NT	NT	NT
	38	10	7.82/8.36	NT	NT	NT
	39	10	7.88/7.84	NT	NT	NT
25	40	83	7.94/7.61	NT	NT	NT
	41	3700	5.68/5.96	NT	NT	NT
	42	370	6.56/6.26	NT	NT	NT
	43	19	8.97/8.61	NT	NT	NT
	44	16	8.23/7.70	NT	NT	NT
30	45	4.4	8.41/8.24	NT	NT	NT
	46	110	6.80/6.64	NT	NT	NT

178

Test	<sup>1</sup> Assay A		<sup>2</sup> Assay B	<sup>3</sup> Assay C		
	Compound	IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition	Duration
	Example #	(nM)		(mg/kg)	(%)	(min.)
5	47	21	7.85/7.58	NT	NT	NT
	48	680	6.27/6.75	NT	NT	NT
	49	120	7.06/7.07	NT	NT	NT
	50	54	7.71/7.89	NT	NT	NT
	51	8.7	8.39/8.51	NT	NT	NT
10	52	100	8.14/8.12	NT	NT	NT
	53	65	7.56/7.83	NT	NT	NT
	54	3100	6.02	NT	NT	NT
	55	80	6.56/7.13	NT	NT	NT
	56	5.0	9.04/8.35	NT	NT	NT
15	57	2300	6.00	NT	NT	NT
	58	140	6.45/6.57	NT	NT	NT
	59	120	7.23/7.59	NT	NT	NT
	60	2200	6.40/6.03	NT	NT	NT
	61	110	7.29/7.70	NT	NT	NT
20	62	26	8.69/8.61	NT	NT	NT
	63	61	7.77/7.67	NT	NT	NT
	64	54	7.00/6.77	NT	NT	NT
	65	23	7.85/7.75	NT	NT	NT
	66	12	9.34/8.58	NT	NT	NT
25	67	3100	5.88/5.78	NT	NT	NT
	68	8.6	8.19/8.65	NT	NT	NT
	69	15	7.80/8.28	NT	NT	NT
	70	44	7.71/8.05	NT	NT	NT
	71	12,000	*	NT	NT	NT
30	72	83	6.11/6.10	NT	NT	NT
	73	790	7.65/7.46	NT	NT	NT

179

Test Compound Example #	<sup>1</sup> Assay A		<sup>2</sup> Assay B	<sup>3</sup> Assay C		
	IC <sub>50</sub>		pA <sub>2</sub>	Dose	Inhibition	Duration
	(nM)			(mg/kg)	(%)	(min.)
5	74	6.5	8.56/8.39	NT	NT	NT
	75	570	6.00/5.45	NT	NT	NT
	76	5400	5.52/5.78	NT	NT	NT
	77	15,000	5.77	NT	NT	NT
	78	101	7.0		93	60-100
10	79	4.9	9.2		100	>200
					50	>180
	80	25	8.1		NT	NT
	81	18	8.0		40	180
	82	7.9	8.5		20	180
15	83	3.6	8.3		15	>180
	84	16	7.1		20	30
	85	8.7	8.9		NT	NT
	86	9	7.8		NT	NT
	87	91	7.8		NT	NT
20	88	50	7.7		NT	NT
	89	18	7.9		NT	NT
	90	5.6	9.0		NT	NT
	91	30	8.6		40	>180
	92	35	7.9		NT	NT
25	93	480	NT		NT	NT
	94	5,800	NT		NT	NT
	95	66	8.2		NT	NT
	96	21	8.0		NT	NT
	97	280	7.7		NT	NT
30	98	22	8.1		NT	NT
	99	280	6.5		NT	NT
	100	4.4	9.4		NT	NT
	101	36	7.8		NT	NT

180

Test	<sup>1</sup> Assay A		<sup>2</sup> Assay B	<sup>3</sup> Assay C	
	Compound	IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition
	Example #	(nM)		(mg/kg)	(%)
5	102	43	7.7		NT
	103	12	8.0		NT
	104	15	8.0		NT
	105	290	6.6		NT
	106	48	7.7		NT
10	107	180	8.3		NT
	108	720	5.3	100	45
	109	250	7.3	30	50
	110	590	6.4		NT
	111	45	9.0	30	87
15	112	2000	5.2		NT
	113	12	8.4	10	60
	114	400	6.4		NT
	115	11	8.2	3	40
	116	230	6.5		NT
20	117	170	6.5		NT
	118	37	9.21/9.17	10	70
	119	16	9.21/9.00	3	20
	120	25	9.05/8.77	10	80
	121	46	NT		NT
25	122	46	NT		NT
	123	50	NT		NT
	124	40	9.42/9.12	3	45
	125	40	9.25/8.80	3	35

181

Test	<sup>1</sup> Assay A	<sup>2</sup> Assay B	<sup>3</sup> Assay C		
Compound	IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition	Duration
Example #	(nM)		(mg/kg)	(%)	(min.)
5	126	240	7.20/7.05	NT	
	127	12,000	4.96	NT	
	128	16	8.63/8.40	NT	
	129	6,700	5.30	NT	
	130	40	8.10/7.94	NT	
10	131	9.5	7.53/8.25		
	132	12	8.6	NT	
	133	10	8.7	3	20
					180
15	134	22	9.3	3	35
	135	16	8.5	3	35
	136	NT	NT		NT
	137	220	8.3		NT
	138	130	8.2		NT
	139	0.270	6.3		NT
	140	0.031	8.1	100	160
	141	0.110	8.02	NT	NT
20	142	2,000	NA	NT	NT
	143	0.052	7.7	85	75
	144	0.088	7.7	50	125
25	145	0.480	6.7	NT	NT
	146	0.072	6.4	NT	NT

182

Test	<sup>1</sup> Assay A	<sup>2</sup> Assay B	<sup>3</sup> Assay C			
Compound	IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition	Duration	
Example #	(nM)		(mg/kg)	(%)	(min.)	
5	147	5.8	5.6	3	74	5-10
	148	0.87	5.8	3	92	20-30
	149	1.1	6.1	3	NT	NT
	150	14	8.03/7.80	3	25	>180
	151	17	7.76/7.97	3	15	180
10	152	150	7.46/7.23	3	10	140
	153	13	8.30/7.69	3	25	>180
	154	97	8.19/8.38		NA	
	155	86	7.60/7.14		NA	
	156	78	8.03/7.66		NA	
15	157	530	- /6.22		NA	
	158	54	8.23/8.14	3	30	>180
	159	21	7.92/7.56	3	10	150
	160	64	7.87/7.71			
	161	28			NA	
20	162	380	6.21/6.55		NA	
	163	420	7.42/6.75		NA	
	164	1700			NA	
	165	410	6.90/7.18		NA	



183

Test Compound Example #	<sup>1</sup> Assay A	<sup>2</sup> Assay B	<sup>3</sup> Assay C		
	IC <sub>50</sub> (nM)	pA <sub>2</sub>	Dose (mg/kg)	Inhibition (%)	Duration (min.)
5	166	160	7.57/7.74		NA
	167	370	7.08/7.11		NA
	168	420	7.69/7.58		NA
	169	150	7.78/7.58	3	15
	170	26	7.08/7.77	3	40
10	171	28	7.52/7.11	3	0
	172	70	7.15/7.04		NA
	173	90	7.49/6.92		NA
	174	180	7.29/7.02		NA
	175	27	NA	3	0
15	176	9.8	7.69/7.55	3	10
	177	26	7.41/7.85	3	15
	178	88	7.54/7.47		NA
	179	310	6.67/ -		NA
	180	26	7.56/7.15	3	25
20	181	21	7.70/7.12	3	20
	182	59	NA		NA
	183	390	NA		NA
	184	1100	6.78/ -		NA

184

Test	<sup>1</sup> Assay A		<sup>2</sup> Assay B	<sup>3</sup> Assay C		
	Compound	IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition	Duration
	Example #	(nM)		(mg/kg)	(%)	(min.)
5	185	6.5	8.82/8.53	3	50	> 180
	186	38	8.13/7.40	3	25	180
	187	770	7.46/6.95		NA	
	188	140	7.72/7.09		NA	
	189	29	8.64/8.23		NA	
10	190	10	7.87/7.89	3	10	180
	191	81	7.75/7.76	3	10	180
	192	140			NA	
	193	11	9.27/8.87	3	10	180
	194	47	7.64/7.35		NA	
15	195	34	8.44/8.03		NA	
	196	31	7.68/8.26		NA	
	197	14	8.03/8.60		NA	
	198	7.6	8.76/8.64	3	35	> 180
	199	10	8.79/8.85	3	60	> 180
20	200	20	8.42/8.77	3	45	> 180
	201	17	8.78/8.63	3	10	180
	202	12	8.79/8.64	3	65	> 180
	203	9.2	8.43/8.36	3	50	> 180
	204	16	9.17/8.86	3	75	> 180
25	205	20	9.14/9.15	3	40	> 180
	206	5.4	8.75/8.89	3	30	> 180
	207	99	9.04/8.60		NA	
	208	22	9.19/8.69	3	50	> 180
	209	5.0	9.41/9.16	3	25	> 180
30	210	3.6	8.36/8.44	3	15	180
	211	18	8.74/8.67	3	35	> 180
	212	23	8.85/8.25	3	15	180
	213	51	NA		NA	
	214	65	NA		NA	
35	215	45	NA		NA	
	216	5.4	8.80/9.04	3	50	> 180

185

Test	<sup>1</sup> Assay A	<sup>2</sup> Assay B	<sup>3</sup> Assay C		
Compound	IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition	Duration
Example #	(nM)		(mg/kg)	(%)	(min.)
5					
217	9.4	NA	3	65	> 180
218	9.0	NA		NA	
219	14	NA		NA	
220	7.0	NA	3	75	120
10 221	4.8	NA	3	25	> 180
222	5.0	NA		NA	
223	14	7.45/7.87	3	20	> 180
224	91	NA		NA	
225	160	NA		NA	
15 226	93	NA		NA	
227	89	7.55/7.67		NA	
228	4.5	9.17/8.25	3	80	>180
229	19	NT	3	40	>180
230	2.6	8.23/8.69	3	25	>180
20 231	3.6	NT	3	75	>180
232	4.4	8.59/8.89	3	70	>180
233	84	8.51/8.78		NT	
234	5.0	8.49/9.00	3	20	-
235	34	7.14/7.07		NT	
25 236	4.9	NC	3	70	>180
237	3.6	NT		NT	
238	1.7	NT	3	15	>180
239	6.8	7.88/8.01	3	20	>180
240	120	NA		NA	
30 241	6.9	8.57/8.24	3	40	>180
242	110	7.11/6.60		NA	
243	250	NA		NA	
244	150	7.17/7.17		NA	
245	98	6.64/7.04		NA	
35 246	72	7.46/7.59		NA	
247	9.4	8.26/8.41	3	20	180

186

Test	<sup>1</sup> Assay A	<sup>2</sup> Assay B	<sup>3</sup> Assay C		
Compound	IC <sub>50</sub>	pA <sub>2</sub>	Dose	Inhibition	Duration
Example #	(nM)		(mg/kg)	(%)	(min.)
5	248	20	7.68/7.50	3	10
	249	4.4	NA	3	20
	250	43	NA	3	0
	251	25	NA		NA
	252	13	NA		NA
10	253	2.0	NA		NA
	254	72	NA		NA
	255	12	7.61/7.46	3	20
	256	4.1	8.43/7.78	3	30
	257	160	6.63/6.68		NA
15	258	350	6.84/6.84		NA
	259	54	NA		NA
	260	220	NA		NA
	261	18	NA		NA
	262	530	-/6.22		NA
20	263	57	NA		NA
	264	11	NA		NA
	265	110	NA		NA
	266	290	NA		NA
	267	25	NA	3	25
25	268	520	NA	3	0
	269	9.7	NA		NA
	270	21	NA		NA
	271	14	NC	3	20%
	272	97	NC	3	70%
30	273	9.8	8.53/8.61	3	25%
	274	13	9.06/8.85	3	35%
	275	6.3	9.07/ --	3	40%
	276	33	8.71/8.64	3	<20%
	277	190	-- /6.54		NT
35	278	30	8.49/8.51	3	50%
	279	270	8.06/8.25		NT
	280	480	6.41/6.35	NT	NT

187

NT = NOT TESTED

NC = Non-Competitive antagonist

5 \*Antagonist Activity not observed up to 10  $\mu$ M of test compound.

1 Assay A: Angiotensin II Binding Activity

2 Assay B: In Vitro Vascular Smooth Muscle Response

10 3 Assay C: In Vivo Pressor Response

15 Test Compounds administered intragastrically, except for compounds of examples #1-#2, #4-#25, #27-#29, #30-#79, #108-#109, #111, #118 and #139-#149 which were given intraduodenally.

Administration of the angiotensin II receptor antagonist and the aldosterone receptor antagonist may take place sequentially in separate formulations, or may be accomplished by simultaneous administration in a single formulation or separate formulations. Administration may be accomplished by oral route, or by intravenous, intramuscular or subcutaneous injections. The formulation may be in the form of a bolus, or in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more pharmaceutically-acceptable carriers or diluents, or a binder such as gelatin or hydroxypropyl-methyl cellulose, together with one or more of a lubricant, preservative, surface-active or dispersing agent.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. These may with advantage contain an amount of each active ingredient from about 1 to 250 mg, preferably from about 25 to 150 mg. A suitable daily dose for a mammal may vary widely depending on the condition of the patient and other factors. However, a dose of from about 0.01 to 30 mg/kg body weight, particularly from about 1 to 15 mg/kg body weight, may be appropriate.

The active ingredients may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier. A suitable daily dose of each active component is from about 0.01 to 15 mg/kg body weight injected per day in multiple doses depending on the disease being treated. A preferred

daily dose would be from about 1 to 10 mg/kg body weight. Compounds indicated for prophylactic therapy will preferably be administered in a daily dose generally in a range from about 0.1 mg to about 15 mg per kilogram of body weight per day. A more preferred dosage will be a range from about 1 mg to about 15 mg per kilogram of body weight. Most preferred is a dosage in a range from about 1 to about 10 mg per kilogram of body weight per day. A suitable dose can be administered, in multiple sub-doses per day. These sub-doses may be administered in unit dosage forms. Typically, a dose or sub-dose may contain from about 1 mg to about 100 mg of active compound per unit dosage form. A more preferred dosage will contain from about 2 mg to about 50 mg of active compound per unit dosage form. Most preferred is a dosage form containing from about 3 mg to about 25 mg of active compound per unit dose.

In combination therapy, the aldosterone receptor antagonist may be present in an amount in a range from about 5 mg to about 400 mg, and the AII antagonist may be present in an amount in a range from about 1 mg to about 800 mg, which represents aldosterone antagonist-to-AII antagonist ratios ranging from about 400:1 to about 1:160.

In a preferred combination therapy, the aldosterone receptor antagonist may be present in an amount in a range from about 10 mg to about 200 mg, and the AII antagonist may be present in an amount in a range from about 5 mg to about 600 mg, which represents aldosterone antagonist-to-AII antagonist ratios ranging from about 40:1 to about 1:60.

In a more preferred combination therapy, the aldosterone receptor antagonist may be present in an amount in a range from about 20 mg to about 100 mg, and the AII antagonist may be present in an amount in a range from about 10 mg to about 400 mg, which represents aldosterone

antagonist-to-AII antagonist ratios ranging from about 10:1 to about 1:20.

The dosage regimen for treating a disease  
5 condition with the combination therapy of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex and medical condition of the patient, the severity of the disease, the route of administration, and the particular compound employed, and  
10 thus may vary widely.

For therapeutic purposes, the active components of this combination therapy invention are ordinarily combined with one or more adjuvants appropriate to the indicated  
15 route of administration. If administered per os, the components may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric  
20 acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active  
25 compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the  
30 carriers or diluents mentioned for use in the formulations for oral administration. The components may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and  
35 modes of administration are well and widely known in the pharmaceutical art.



Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

What Is Claimed Is:

1. A combination comprising a therapeutically-effective amount of an angiotensin II receptor antagonist  
5 and a therapeutically-effective non-diuretic-effective amount of an epoxy-free spiro lactone-type aldosterone receptor antagonist.

2. The combination of Claim 1 wherein said  
10 aldosterone receptor antagonist is selected from spiro lactone-type compounds of Formula A

15 (A)

wherein R is lower alkyl of up to 5 carbon  
atoms, and

20

25

3. The combination of Claim 2 wherein said spiro lactone-type compound is selected from compounds of the group consisting of:

30 7 $\alpha$ -Acetylthio-3-oxo-4,15-androstadiene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
3-Oxo-7 $\alpha$ -propionylthio-4,15-androstadiene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
6 $\beta$ ,7 $\beta$ -Methylene-3-oxo-4,15-androstadiene-[17( $\beta$ -1')-  
35 spiro-5']perhydrofuran-2'-one;  
15 $\alpha$ ,16 $\alpha$ -Methylene-3-oxo-4,7 $\alpha$ -propionylthio-4-androstene[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;

6 $\beta$ , 7 $\beta$ , 15 $\alpha$ , 16 $\alpha$ -Dimethylene-3-oxo-4-androstene  
[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
7 $\alpha$ -Acetylthio-15 $\beta$ , 16 $\beta$ -Methylene-3-oxo-4-androstene-  
[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one;  
5 15 $\beta$ , 16 $\beta$ -Methylene-3-oxo-7 $\beta$ -propionylthio-4-  
androstene-[17( $\beta$ -1')-spiro-5']perhydrofuran-2'-one; and  
6 $\beta$ , 7 $\beta$ , 15 $\beta$ , 16 $\beta$ -Dimethylene-3-oxo-4-androstene-[17( $\beta$ -  
1')-spiro-5']perhydrofuran-2'-one.

10 4. The combination of Claim 1 wherein said  
aldosterone receptor antagonist is selected from  
spiro lactone-type compounds of Formula B:

15

(B)

wherein

20

R<sup>1</sup> is C<sub>1-3</sub>-alkyl or C<sub>1-3</sub> acyl and R<sup>2</sup> is hydrogen or C<sub>1-3</sub>-  
alkyl.

25 5. The combination of Claim 4 wherein said  
spiro lactone-type compound is selected from:

1 $\alpha$ -Acetylthio-15 $\beta$ , 16 $\beta$ -methylene-7 $\alpha$ -methylthio-3-oxo-  
17 $\alpha$ -pregn-4-ene-21, 17-carbolactone; and  
15 $\beta$ , 16 $\beta$ -Methylene-1 $\alpha$ , 7 $\alpha$ -dimethylthio-3-oxo-17 $\alpha$ -  
30 pregn-4-ene-21, 17-carbolactone.

6. The combination of Claim 1 wherein said aldosterone receptor antagonist is selected from spirolactone-type compounds of Formula C:

5

(C)

10

7. The combination of Claim 6 wherein said spirolactone-type compound is selected from

7 $\alpha$ -Acylthio-21-hydroxy-3-oxo-17 $\alpha$ -pregn-4-ene-17-carboxylic acid lactone;

15

21-hydroxy-3-oxo-17 $\alpha$ -pregn-1,4-diene-17-carboxylic acid lactone; and

17-hydroxy-7 $\alpha$ -mercapto-3-oxo-17 $\alpha$ -pregn-4-ene-21-carboxylic acid  $\gamma$ -lactone acetate.

20

8. The combination of Claim 1 wherein said angiotensin II receptor antagonist is selected from compounds consisting of a first portion and a second portion, wherein said first portion is selected from a fragment of Formula I:

25

Ar-Alk-L

Ar-L-Ar-Alk-L

30

Het-L-Ar-Alk-L

Het-L-Het-Alk-L

(I)

Ar-L-Het-Alk-L

Het-L-Alk-L

35



wherein Ar is a five or six-membered carbocyclic ring system consisting of one ring or two fused rings, with such ring or rings being fully unsaturated or partially or fully saturated;

wherein Het is a monocyclic or bicyclic fused ring system having from five to eleven ring members, and having at least one of such ring members being a hetero atom selected from one or more hetero atoms selected from oxygen, nitrogen and sulfur, and with such ring system containing up to six of such hetero atoms as ring members;

wherein Alk is an alkyl radical or alkylene chain, linear or branched, containing from one to about five carbon atoms;

wherein L is a straight bond or a bivalent linker moiety selected from carbon, oxygen and sulfur;

and wherein said second portion is a monocyclic heterocyclic moiety selected from moieties of Formula IIa or is a bicyclic heterocyclic moiety selected from moieties of Formula IIb:



wherein each of X¹ through X⁶ is selected from -CH=, -CH₂-, -N=, -NH-, O, and S, with the proviso that at least one of X¹ through X⁶ in each of Formula IIa and Formula IIb must be a hetero atom, and wherein said

heterocyclic moiety of Formula IIa or IIb may be attached through a bond from any ring member of the Formula IIa or IIb heterocyclic moiety having a substitutable or a bond-forming position.

5

9. The combination of Claim 8 wherein said monocyclic heterocyclic moiety of Formula IIa is selected from thienyl, furyl, pyranyl, pyrrolyl, imidazolyl, triazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, 10 pyridazinyl, isothiazolyl, isoxazolyl, furazanyl, pyrrolidinyl, pyrrolinyl, furanyl, thiophenyl, isopyrrolyl, 3-isopyrrolyl, 2-isoimidazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2-dithiolyl, 1,3-dithiolyl, 1,2,3-oxathiolyl, oxazolyl, thiazolyl, 1,2,3-oxadiazolyl, 15 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3,4-oxatriazolyl, 1,2,3,5-oxatriazolyl, 1,2,3-dioxazolyl, 1,2,4-dioxazolyl, 1,3,2-dioxazolyl, 1,3,4-dioxazolyl, 1,2,5-oxathiazolyl, 1,3-oxathiolyl, 1,2-pyranyl, 1,4-pyranyl, 1,2-pyronyl, 1,4-pyronyl, 20 pyridinyl, piperazinyl, s-triazinyl, as-triazinyl, v-triazinyl, 1,2,4-oxazinyl, 1,3,2-oxazinyl, 1,3,6-oxazinyl, 1,2,6-oxazinyl, 1,4-oxazinyl, o-isoxazinyl, p-isoxazinyl, 1,2,5-oxathiazinyl, 1,4-oxazinyl, o-isoxazinyl, p-isoxazinyl, 1,2,5-oxathiazinyl, 1,2,6-oxathiazinyl, 25 oxathiazinyl, 1,4,2-oxadiazinyl, 1,3,5,2-oxadiazinyl, morpholinyl, azepinyl, oxepinyl, thiepinyl and 1,2,4-diazepinyl.

10. The combination of Claim 9 wherein said 30 bicyclic heterocyclic moiety of Formula IIb is selected from benzo[b]thienyl, isobenzofuranyl, chromenyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolizinyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, 35 pteridinyl, isochromanyl, chromanyl, thieno[2,3-b]furanyl, 2H-furo[3,2-b]pyranyl, 5H-pyrido[2,3-d][1,2]oxazinyl, 1H-pyrazolo[4,3-d]oxazolyl, 4H-

imidazo[4,5-d]thiazolyl, pyrazino[2,3-d]pyridazinyl,  
imidazo[2,1-b]thiazolyl, cyclopenta[b]pyranyl, 4H-  
[1,3]oxathiololo-[5,4-b]pyrrolyl, thieno[2,3-b]furanyl,  
imidazo[1,2-b][1,2,4]triazinyl and  
5 4H-1,3-dioxolo[4,5-d]imidazolyl.

11. The combination of Claim 10 wherein said  
angiotensin II receptor antagonist compound having said  
first-and-second-portion moieties of Formula I and II is  
10 further characterized by having an acidic moiety attached  
to either of said first-and-second-portion moieties.

12. The combination of Claim 11 wherein said  
acidic moiety is attached to the first-portion moiety of  
15 Formula I and is defined by Formula III:




wherein n is a number selected from zero through three,  
20 inclusive, and wherein A is an acidic group selected to  
contain at least one acidic hydrogen atom, and the amide,  
ester and salt derivatives of said acidic moieties;  
wherein U is a spacer group independently selected from  
one or more of alkyl, cycloalkyl, cycloalkylalkyl,  
25 alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one  
or more ring atoms selected from oxygen, sulfur and  
nitrogen atoms.


13. The combination of Claim 12 wherein said  
30 acidic moiety is selected from carboxyl moiety and  
tetrazolyl moiety.

14. The combination of Claim 12 wherein any of  
the moieties of Formula I and Formula II may be  
35 substituted at any substitutable position by one or more  
radicals selected from hydrido, hydroxy, alkyl, alkenyl,  
alkynyl, aralkyl, hydroxyalkyl, haloalkyl, halo, oxo,

- alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxy carbonyloxy, alkylcarbonyl, alkoxy carbonyl, 5 aralkoxy carbonyl, carboxyl, mercapto, mercapto carbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, 10 sulfur and nitrogen atoms, and amino and amido radicals of the formula




- 15 wherein W is oxygen atom or sulfur atom; wherein each of R<sup>1</sup> through R<sup>5</sup> is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, YR<sup>6</sup> and



- 20 wherein Y is selected from oxygen atom and sulfur atom and R<sup>6</sup> is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup> and R<sup>8</sup> is independently selected from 25 hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of 30 R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup> and R<sup>8</sup> is further independently selected from amino and amido radicals of the formula



199



wherein W is oxygen atom or sulfur atom;  
 wherein each of R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is  
 5 independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R<sup>2</sup> and R<sup>3</sup> taken together and each of R<sup>4</sup> and R<sup>5</sup> taken together may form a heterocyclic group  
 10 having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or  
 15 partially unsaturated; wherein each of R<sup>2</sup> and R<sup>3</sup> taken together and each of R<sup>7</sup> and R<sup>8</sup> taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further  
 20 contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

15. The combination of Claim 14 wherein said  
 25 angiotensin II receptor antagonist is 5-[2-[5-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl]-2-pyridinyl]phenyl-1H-tetrazole or a pharmaceutically-acceptable salt thereof and said spiro lactone-type aldosterone receptor antagonist is  
 30 17-hydroxy-7 $\alpha$ -mercapto-3-oxo-17 $\alpha$ -pregn-4-ene-21-carboxylic acid  $\gamma$ -lactone acetate or a pharmaceutically-acceptable salt thereof.

16. The combination of Claim 15 further

200

characterized by said angiotensin II receptor antagonist and said aldosterone receptor antagonist being present in said combination in a weight ratio range from about one-to-one to about twenty-to-one of said angiotensin II  
5 receptor antagonist to said aldosterone receptor antagonist.

17. The combination of Claim 15 wherein said weight ratio range is from about five-to-one to about  
10 fifteen-to-one.

18. The combination of Claim 17 wherein said weight ratio range is about ten-to-one.

15 19. The combination of Claim 1 wherein said angiotensin II receptor antagonist is selected from the group consisting of:  
saralasin acetate, candesartan cilexetil, CGP-63170, EMD-66397, KT3-671, LR-B/081, valsartan, A-81282,  
20 BIBR-363, BIBS-222, BMS-184698, candesartan, CV-11194, EXP-3174, KW-3433, L-161177, L-162154, LR-B/057, LY-235656, PD-150304, U-96849, U-97018, UP-275-22, WAY-126227, WK-1492.2K, YM-31472, losartan potassium, E-4177, EMD-73495, eprosartan, HN-65021, irbesartan,  
25 L-159282, ME-3221, SL-91.0102, Tasosartan, Telmisartan, UP-269-6, YM-358, CGP-49870, GA-0056, L-159689, L-162234, L-162441, L-163007, PD-123177, A-81988, BMS-180560, CGP-38560A, CGP-48369, DA-2079, DE-3489, DuP-167, EXP-063, EXP-6155, EXP-6803, EXP-7711, EXP-9270, FK-739,  
30 HR-720, ICI-D6888, ICI-D7155, ICI-D8731, isoteoline, KRI-1177, L-158809, L-158978, L-159874, LR B087, LY-285434, LY-302289, LY-315995, RG-13647, RWJ-38970, RWJ-46458, S-8307, S-8308, saprisartan, saralasin, Sarmesin, WK-1360, X-6803, ZD-6888, ZD-7155, ZD-8731,  
35 BIBS39, CI-996, DMP-811, DuP-532, EXP-929, L-163017, LY-301875, XH-148, XR-510, zolasartan and PD-123319.

20. The combination of Claim 19 wherein said angiotensin II receptor antagonist is selected from the group consisting of:

saralasin acetate, candesartan cilexetil, CGP-63170,  
5 EMD-66397, KT3-671, LR-B/081, valsartan, A-81282,  
BIBR-363, BIBS-222, BMS-184698, candesartan, CV-11194,  
EXP-3174, KW-3433, L-161177, L-162154, LR-B/057,  
LY-235656, PD-150304, U-96849, U-97018, UP-275-22,  
WAY-126227, WK-1492.2K, YM-31472, losartan potassium,  
10 E-4177, EMD-73495, eprosartan, HN-65021, irbesartan,  
L-159282, ME-3221, SL-91.0102, Tasosartan, Telmisartan,  
UP-269-6, YM-358, CGP-49870, GA-0056, L-159689, L-162234,  
L-162441, L-163007 and PD-123177.

15 21. A co-therapy for treating cardiovascular disorders in a subject afflicted with or susceptible to multiple cardiovascular disorders, wherein said co-therapy comprises administering a therapeutically-effective amount of an angiotensin II receptor antagonist  
20 and administering a therapeutically effective non-diuretic-effective amount of an epoxy-free spiro lactone-type aldosterone receptor antagonist.

22. The co-therapy of Claim 21 wherein said  
25 subject is afflicted with or susceptible to or afflicted with hypertension.

23. The co-therapy of Claim 21 wherein said  
subject is susceptible to or afflicted with congestive  
30 heart failure.

24. The co-therapy of Claim 21 further characterized by administering said angiotensin II receptor antagonist and said aldosterone receptor  
35 antagonist in a sequential manner.

25. The co-therapy of Claim 21 further

characterized by administering said angiotensin II receptor antagonist and said aldosterone receptor antagonist in a substantially simultaneous manner.

5           26. The co-therapy of Claim 21 wherein said angiotensin II receptor antagonist is 5-[2-[5-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl]-2-pyridinyl]phenyl-1H-tetrazole or a pharmaceutically-acceptable salt thereof and said aldosterone receptor antagonist is  
10 17-hydroxy-7 $\alpha$ -mercapto-3-oxo-17 $\alpha$ -pregn-4-ene-21-carboxylic acid  $\gamma$ -lactone acetate or a pharmaceutically-acceptable salt thereof.

          27. The co-therapy of Claim 25 further  
15 characterized in administering said angiotensin II receptor antagonist and said aldosterone receptor antagonist is a weight ratio range from about two-to-one to about fifty-to-one of said angiotensin II receptor antagonist to said aldosterone receptor antagonist.

20

          28. The co-therapy of Claim 27 wherein said weight ratio range is from about two-to-one to about ten-to-one.

25           29. The co-therapy of Claim 28 wherein said weight ratio range is about five-to-one.

          30. A method to treat a subject susceptible to or afflicted with congestive heart failure, which method  
30 comprises administering a combination of drug agents comprising a therapeutically-effective amount of an angiotensin II receptor antagonist and a therapeutically-effective non-diuretic-effective amount of an epoxy-free spiro lactone-type aldosterone receptor antagonist.


35

          31. The method of Claim 30 wherein said aldosterone receptor antagonist is 17-hydroxy-7 $\alpha$ -

mercapto-3-oxo-17 $\alpha$ -pregn-4-ene-21-carboxylic acid  $\gamma$ -lactone acetate or a pharmaceutically-acceptable salt thereof.

5           32. The method of Claim 30 wherein said angiotensin II receptor antagonist is selected from compounds consisting of a first portion and a second portion, wherein said first portion is selected from a fragment of Formula I:

10



15

wherein Ar is a five or six-membered carbocyclic ring system consisting of one ring or two fused rings, with such ring or rings being fully unsaturated or partially or fully saturated;

20

wherein Het is a monocyclic or bicyclic fused ring system having from five to eleven ring members, and having at least one of such ring members being a hetero atom selected from one or more hetero atoms selected from oxygen, nitrogen and sulfur, and with such ring system containing up to six of such hetero atoms as ring members;

30



wherein Alk is an alkyl radical or alkylene chain, linear or branched, containing from one to about five carbon atoms;

35

wherein L is a straight bond or a bivalent linker moiety selected from carbon, oxygen and sulfur;

and wherein said second portion is a monocyclic heterocyclic moiety selected from moieties of Formula IIa or is a bicyclic heterocyclic moiety selected from moieties of Formula IIb:

5



wherein each of X<sup>1</sup> through X<sup>6</sup> is selected from -CH=,  
 10 -CH<sub>2</sub>-, -N=, -NH-, O, and S, with the proviso that at least one of X<sup>1</sup> through X<sup>6</sup> in each of Formula IIa and Formula IIb must be a hetero atom, and wherein said heterocyclic moiety of Formula IIa or IIb may be attached through a bond from any ring member of the Formula IIa or  
 15 IIb heterocyclic moiety having a substitutable or a bond-forming position.

33. The method of Claim 32 wherein said monocyclic heterocyclic moiety of Formula IIa is selected  
 20 from thienyl, furyl, pyranyl, pyrrolyl, imidazolyl, triazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, isothiazolyl, isoxazolyl, furazanyl, pyrrolidinyl, pyrrolinyl, furanyl, thiophenyl, isopyrrolyl, 3-isopyrrolyl, 2-isoimidazolyl, 1,2,3-  
 25 triazolyl, 1,2,4-triazolyl, 1,2-dithiolyl, 1,3-dithiolyl, 1,2,3-oxathiolyl, oxazolyl, thiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3,4-oxatriazolyl, 1,2,3,5-oxatriazolyl, 1,2,3-

dioxazolyl, 1,2,4-dioxazolyl, 1,3,2-dioxazolyl, 1,3,4-dioxazolyl, 1,2,5-oxathiazolyl, 1,3-oxathiolyl, 1,2-pyranyl, 1,4-pyranyl, 1,2-pyranyl, 1,4-pyranyl, pyridinyl, piperazinyl, s-triazinyl, as-triazinyl, v-triazinyl, 1,2,4-oxazinyl, 1,3,2-oxazinyl, 1,3,6-oxazinyl, 1,2,6-oxazinyl, 1,4-oxazinyl, o-isoxazinyl, p-isoxazinyl, 1,2,5-oxathiazinyl, 1,4-oxazinyl, o-isoxazinyl, p-isoxazinyl, 1,2,5-oxathiazinyl, 1,2,6-oxathiazinyl, 1,4,2-oxadiazinyl, 1,3,5,2-oxadiazinyl, morpholinyl, azepinyl, oxepinyl, thiepinyl and 1,2,4-diazepinyl.

34. The method of Claim 33 wherein said bicyclic heterocyclic moiety of Formula IIb is selected from benzo[b]thienyl, isobenzofuranyl, chromenyl, indolizinyl, isoindolyl, indolyl, indazolyl, purinyl, quinolizinyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, isochromanyl, chromanyl, thieno[2,3-b]furanyl, 2H-furo[3,2-b]pyranyl, 5H-pyrido[2,3-d][1,2]oxazinyl, 1H-pyrazolo[4,3-d]oxazolyl, 4H-imidazo[4,5-d]thiazolyl, pyrazino[2,3-d]pyridazinyl, imidazo[2,1-b]thiazolyl, cyclopenta[b]pyranyl, 4H-[1,3]oxathiolo-[5,4-b]pyrrolyl, thieno[2,3-b]furanyl, imidazo[1,2-b][1,2,4]triazinyl and 4H-1,3-dioxolo[4,5-d]imidazolyl.

35. The method of Claim 34 wherein said angiotensin II receptor antagonist compound having said first-and-second-portion moieties of Formula I and II is further characterized by having an acidic moiety attached to either of said first-and-second-portion moieties.

36. The method of Claim 35 wherein said acidic moiety is attached to the first-portion moiety of Formula I and is defined by Formula III:

206


-U<sub>n</sub>A

(III)

wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein U is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms.

37. The method of Claim 36 wherein said acidic moiety is selected from carboxyl moiety and tetrazolyl moiety.


38. The method of Claim 36 wherein any of the moieties of Formula I and Formula II may be substituted at any substitutable position by one or more radicals selected from hydrido, hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, haloalkyl, halo, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxy carbonyloxy, alkylcarbonyl, alkoxy carbonyl, aralkoxy carbonyl, carboxyl, mercapto, mercaptocarbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino and amido radicals of the formula






207

wherein W is oxygen atom or sulfur atom; wherein each of R<sup>1</sup> through R<sup>5</sup> is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, YR<sup>6</sup> and



5

wherein Y is selected from oxygen atom and sulfur atom and R<sup>6</sup> is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup> and R<sup>8</sup> is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup> and R<sup>8</sup> is further independently selected from amino and amido radicals of the formula



20

wherein W is oxygen atom or sulfur atom; wherein each of R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R<sup>2</sup> and R<sup>3</sup> taken together and each of R<sup>4</sup> and R<sup>5</sup> taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or

30

partially unsaturated; wherein each of R<sup>2</sup> and R<sup>3</sup> taken together and each of R<sup>7</sup> and R<sup>8</sup> taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

10                   39. The method of Claim 38 wherein said angiotensin II receptor antagonist is 5-{2-[5-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl]-2-pyridinyl]phenyl-1H-tetrazole or a pharmaceutically-acceptable salt thereof and said aldosterone receptor antagonist is  
15 17-hydroxy-7 $\alpha$ -mercapto-3-oxo-17 $\alpha$ -pregn-4-ene-21-carboxylic acid  $\gamma$ -lactone acetate or a pharmaceutically-acceptable salt thereof.

                  40. The method of Claim 39 further  
20 characterized by said angiotensin II receptor antagonist and said aldosterone receptor antagonist being present in said combination in a weight ratio range from about one-to-one to about twenty-to-one of said angiotensin II receptor antagonist to said aldosterone receptor  
25 antagonist.

                  41. The method of Claim 40 wherein said weight ratio range is from about five-to-one to about fifteen-to-one.

30

                  42. The method of Claim 41 wherein said weight ratio range is about ten-to-one.

                  43. The method of Claim 30 wherein said  
35 angiotensin II receptor antagonist is selected from the group consisting of saralasin acetate, candesartan cilexetil, CGP-63170,

- EMD-66397, KT3-671, LR-B/081, valsartan, A-81282,  
BIBR-363, BIBS-222, BMS-184698, candesartan, CV-11194,  
EXP-3174, KW-3433, L-161177, L-162154, LR-B/057,  
LY-235656, PD-150304, U-96849, U-97018, UP-275-22,  
5 WAY-126227, WK-1492.2K, YM-31472, losartan potassium,  
E-4177, EMD-73495, eprosartan, HN-65021, irbesartan,  
L-159282, ME-3221, SL-91.0102, Tasosartan, Telmisartan,  
UP-269-6, YM-358, CGP-49870, GA-0056, L-159689, L-162234,  
L-162441, L-163007, PD-123177, A-81988, BMS-180560,  
10 CGP-38560A, CGP-48369, DA-2079, DE-3489, DuP-167,  
EXP-063, EXP-6155, EXP-6803, EXP-7711, EXP-9270, FK-739,  
HR-720, ICI-D6888, ICI-D7155, ICI-D8731, isoteoline,  
KRI-1177, L-158809, L-158978, L-159874, LR B087,  
LY-285434, LY-302289, LY-315995, RG-13647, RWJ-38970,  
15 RWJ-46458, S-8307, S-8308, saprisartan, saralasin,  
Sarmesin, WK-1360, X-6803, ZD-6888, ZD-7155, ZD-8731,  
BIBS39, CI-996, DMP-811, DuP-532, EXP-929, L-163017,  
LY-301875, XH-148, XR-510, zolasartan and PD-123319.

- 20 44. The method of Claim 43 wherein said  
angiotensin II receptor antagonist is selected from the  
group consisting of  
saralasin acetate, candesartan cilexetil, CGP-63170,  
EMD-66397, KT3-671, LR-B/081, valsartan, A-81282,  
25 BIBR-363, BIBS-222, BMS-184698, candesartan, CV-11194,  
EXP-3174, KW-3433, L-161177, L-162154, LR-B/057,  
LY-235656, PD-150304, U-96849, U-97018, UP-275-22,  
WAY-126227, WK-1492.2K, YM-31472, losartan potassium,  
E-4177, EMD-73495, eprosartan, HN-65021, irbesartan,  
30 L-159282, ME-3221, SL-91.0102, Tasosartan, Telmisartan,  
UP-269-6, YM-358, CGP-49870, GA-0056, L-159689, L-162234,  
L-162441, L-163007 and PD-123177.

**THIS PAGE BLANK (USPTO)**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 45/06, 31/585, 31/41 // (A61K 45/06, 31:585), (A61K 45/06, 31:41)</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 96/40258</b> <b>(43) International Publication Date:</b> 19 December 1996 (19.12.96)
<b>(21) International Application Number:</b> PCT/US96/09342 <b>(22) International Filing Date:</b> 5 June 1996 (05.06.96) <b>(30) Priority Data:</b> 08/486,089 7 June 1995 (07.06.95) US <b>(60) Parent Application or Grant</b> (63) Related by Continuation US 08/486,089 (CON) Filed on 7 June 1996 (07.06.96) <b>(71) Applicant (for all designated States except US):</b> G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MACLAUGHLAN, Todd, E. [CA/US]; 34128 Sulkey Drive, Grayslake, IL 60093 (US). SCHUH, Joseph, R. [US/US]; 2055 Rurline Drive, St. Louis, MO 63146 (US). <b>(74) Agents:</b> KEANE, J., Timothy et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> <b>(88) Date of publication of the international search report:</b> 23 January 1997 (23.01.97)	
<b>(54) Title:</b> SPIRONOLACTONE AND ANGIOTENSIN II ANTAGONIST COMBINATION THERAPY FOR TREATMENT OF CONGESTIVE HEART FAILURE  <b>(57) Abstract</b>  A combination therapy comprising a therapeutically-effective amount of an epoxy-free spiro lactone-type aldosterone receptor antagonist and a therapeutically-effective amount of an angiotensin II receptor antagonist is described for treatment of circulatory disorders, including cardiovascular disorders such as hypertension, congestive heart failure, cirrhosis and ascites. Preferred angiotensin II receptor antagonists are those compounds having high potency and bioavailability and which are characterized in having an imidazole or triazole moiety attached to a biphenylmethyl or pyridinyl/phenylmethyl moiety. A preferred epoxy-free spiro lactone-type aldosterone receptor antagonist is spironolactone. A preferred combination therapy includes the angiotensin II receptor antagonist 5-[2-[(3,5-dibutyl-1H-1,2,4-triazol-1-yl)methyl]-2-pyridinyl]phenyl-1H-tetrazole and the aldosterone receptor antagonist spironolactone.		

**THIS PAGE BLANK (USPTO)**

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam





## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/09342

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K45/06 A61K31/585 A61K31/41 //(A61K45/06,31:585),  
(A61K45/06,31:41)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y P,A	WO,A,95 15166 (CURATORS OF THE UNIVERSITY OF MISSOURI) 8 June 1995 see page 8-12; claims 1,3-7; example 2	1,7,21, 22,30,31 15-18, 26-29, 39-42
X	WO,A,94 09778 (MERCK & CO ) 11 May 1994	1,7-14, 19-23, 30-38, 43,44
Y	see page 1-2; claims 1-3,6-8,10	1,7,21, 22,30,31
A	see page 6, line 9; figures I-XI	15,26,39

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

26 November 1996

Date of mailing of the international search report

05.12.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Kanbier, D

**THIS PAGE BLANK (USPTO)**

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/09342

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANNALS OF INTERNAL MEDICINE, vol. 87, no. 2, 1987, pages 183-187, XP000610206 G.H. ANDERSON ET AL: "DIURETIC THERAPY AND RESPONSE OF ESSENTIAL HYPERTENSION TO SARALASIN"	1,7-14, 19,21, 22,24, 30-38,43
A	see page 184-185; tables 1,2	15,20, 26,39,44
X	--- EP,A,0 481 448 (SQUIBB & SONS) 22 April 1992	1,7-15, 21-23, 26,30-39
A	see page 11, line 20-45; claims 1,6-8,12,13; examples 12-21	16-20, 27-29, 40-44
X	--- WO,A,91 12001 (MERCK & CO INC) 22 August 1991	1,7-15, 21-23, 26,30-39
A	see page 167	19,20, 43,44
X	--- US,A,5 264 447 (MERCK & CO INC) 23 November 1993	1,7-14, 19-23, 30-38, 43,44
A	see column 3-4; claims 1-3	15,26,39
X	--- EP,A,0 628 313 (TAKEDA CHEMICAL INDUSTRIES) 14 December 1994	1,7-14, 21-23, 30-38
A	see page 2-3 see page 9-10; claims 1-17,19,20	15,26,39
A	--- WO,A,91 15206 (DU PONT DE NEMOURS; MERCK & CO) 17 October 1991 see page 21, line 12-20; claims 1-4,6-8 see page 24, line 7-12 see page 24, line 19-30 see page 26, line 1-6	1,8-14, 16,19-22
A	--- US,A,5 049 565 (MERCK & CO INC) 17 September 1991  see column 3-4 see column 8, line 63-65 -----	1,7-15, 19-23, 26, 30-38, 43,44

**THIS PAGE BLANK (USPTO)**

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/09342

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim(s) 21-29 and 30-44  
is(are) directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:

Please see next page.

3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**THIS PAGE BLANK (USPTO)**

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 96/ 09342

FURTHER INFORMATION CONTINUED FROM: PCT/ISA/210

In view of the large number of compounds, which are defined by the general formula/description, used in the claims: 8-14, 32-38, 19, 20, 43, 44, the search had to be restricted for economic reasons. The search was limited to the compounds for which pharmacological data was given and/or the compounds mentioned in the claims, and to the general idea underlying the application (see Guidelines, part B, chapter III, § 3.6).

A compound cannot be sufficiently characterized by its pharmacological profile or its mechanism of action as it is done in Claim 1, 21, 30 as: "angiotensin II receptor antagonist" and "aldosterone receptor antagonist". The search has been executed based on compounds specifically mentioned in Claims 3, 5, 7, 15, 26, 31, 39 and in the examples.

The content of Claims 2, 4 and 6 is unknown because of the missing formulas A, B and C, respectively. These claims could not be searched at all.

Claims searched incompletely: 8-14, 19, 20, 32-38, 43, 44; 1, 21, 30  
Claims not searched : 2, 4, 6

**THIS PAGE BLANK (USPTO)**



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/09342

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9515166	08-06-95	US-A- 5529992	25-06-96
		AU-A- 1210695	19-06-95
		CA-A- 2177848	08-06-95
		EP-A- 0730458	11-09-96
-----	-----	-----	-----
WO-A-9409778	11-05-94	AU-A- 5449194	24-05-94
-----	-----	-----	-----
EP-A-481448	22-04-92	CA-A- 2053148	17-04-92
		JP-A- 4330065	18-11-92
		US-A- 5470975	28-11-95
-----	-----	-----	-----
WO-A-9112001	22-08-91	CA-A- 2075627	14-08-91
		CA-A- 2075637	14-08-91
		EP-A- 0515535	02-12-92
		EP-A- 0517812	16-12-92
		JP-T- 5503530	10-06-93
		JP-T- 5504969	29-07-93
		US-A- 5449682	12-09-95
		WO-A- 9111999	22-08-91
		US-A- 5264439	23-11-93
-----	-----	-----	-----
US-A-5264447	23-11-93	GB-A- 2270073	02-03-94
-----	-----	-----	-----
EP-A-628313	14-12-94	JP-A- 7053373	28-02-95
-----	-----	-----	-----
WO-A-9115206	17-10-91	NONE	
-----	-----	-----	-----
US-A-5049565	17-09-91	NONE	
-----	-----	-----	-----

**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**